Full wwPDB X-ray Structure Validation Report (i)

Oct 27, 2026 – 03:07 pm GMT

PDB ID	:	30XN / pdb_000030xn
Title	:	Human DTX4 RING-DTC domains
Authors	:	Fowkes, M.M.; Nair, A.; Elkins, J.M.
Deposited on	:	2026-05-15
Resolution	:	2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.2
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	239	3% 79% 17% .
1	B	239	7% 85% 11% . .

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 3561 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called E3 ubiquitin-protein ligase DTX4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	229	Total	C	N	O	S	0	2	0
			1737	1100	301	325	11			
1	B	235	Total	C	N	O	S	0	0	0
			1748	1108	302	327	11			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	381	SER	-	expression tag	UNP Q9Y2E6
A	382	MET	-	expression tag	UNP Q9Y2E6
B	381	SER	-	expression tag	UNP Q9Y2E6
B	382	MET	-	expression tag	UNP Q9Y2E6

- Molecule 2 is ADENOSINE-5-DIPHOSPHORIBOSE (CCD ID: APR) (formula: $C_{15}H_{23}N_5O_{14}P_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			36	15	5	14	2		
2	B	1	Total	C	N	O	P	0	0
			36	15	5	14	2		


- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Zn	0	0
			2	2		
3	B	2	Total	Zn	0	0
			2	2		


3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: E3 ubiquitin-protein ligase DTX4



• Molecule 1: E3 ubiquitin-protein ligase DTX4



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α, β, γ	109.09Å 109.09Å 222.01Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.13 – 2.80 29.13 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.8 (29.13-2.80) 99.8 (29.13-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.71 (at 2.80Å)	Xtriage
Refinement program	REFMAC 5.8.0425	Depositor
R, R_{free}	0.226 , 0.294 0.228 , 0.292	Depositor DCC
R_{free} test set	975 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å^2)	70.3	Xtriage
Anisotropy	0.072	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.32 , 58.8	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	3561	wwPDB-VP
Average B, all atoms (Å^2)	83.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.53% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: APR, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|$ >5	RMSZ	#$\|Z\|$ >5
1	A	0.62	0/1793	1.15	3/2450 (0.1%)
1	B	0.60	0/1798	1.12	3/2456 (0.1%)
All	All	0.61	0/3591	1.14	6/4906 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	515	GLU	CB-CA-C	-8.37	95.16	110.63
1	B	515	GLU	CB-CA-C	-7.70	96.38	110.63
1	A	515	GLU	N-CA-CB	6.66	120.47	110.22
1	B	515	GLU	N-CA-CB	5.79	119.14	110.22
1	B	404	PRO	N-CA-C	5.15	116.09	110.58
1	A	576	VAL	N-CA-CB	-5.08	105.60	112.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1737	0	1628	17	0
1	B	1748	0	1647	18	0
2	A	36	0	21	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	36	0	21	1	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
All	All	3561	0	3317	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:389:THR:N	1:B:390:PRO:CD	2.65	0.60
1:A:495:HIS:N	1:A:496:PRO:HD3	2.21	0.56
1:B:442:HIS:CE1	1:B:467:CYS:HB3	2.42	0.55
1:A:464:CYS:SG	1:A:466:THR:OG1	2.61	0.54
1:B:389:THR:HG23	1:B:392:GLU:HB3	1.91	0.52
1:A:539:LYS:NZ	1:A:598:ASP:OD1	2.40	0.52
1:A:442:HIS:CE1	1:A:467:CYS:HB3	2.45	0.51
1:B:389:THR:HA	1:B:392:GLU:CB	2.42	0.49
1:B:389:THR:N	1:B:390:PRO:HD2	2.25	0.49
1:B:457:ASN:O	1:B:458:LYS:C	2.56	0.49
1:B:467:CYS:O	1:B:468:LYS:HB2	2.14	0.48
1:A:543:VAL:HG21	1:A:597:LEU:HD22	1.96	0.47
1:A:591:TYR:HA	1:A:592:PRO:C	2.41	0.46
1:A:457:ASN:O	1:A:458:LYS:C	2.60	0.45
1:B:437:LEU:O	1:B:438:SER:C	2.59	0.45
1:B:392:GLU:O	1:B:393:VAL:HG22	2.16	0.45
1:B:486:HIS:NE2	1:B:502:ARG:HD2	2.31	0.45
1:A:437:LEU:O	1:A:438:SER:C	2.59	0.45
1:A:562:SER:HB3	1:A:565:THR:OG1	2.16	0.45
1:A:550:ALA:HB3	1:A:556:ILE:HD11	1.98	0.44
1:B:591:TYR:HA	1:B:592:PRO:C	2.42	0.44
1:B:389:THR:HA	1:B:392:GLU:HB3	1.99	0.44
1:A:573:TRP:HB2	2:A:701:APR:O4'	2.18	0.44
1:B:394:LEU:O	1:B:395:LYS:C	2.61	0.44
1:B:389:THR:HA	1:B:392:GLU:HB2	2.00	0.44
1:B:578:HIS:O	2:B:701:APR:H2	2.19	0.42
1:A:492:LEU:HD22	1:A:591:TYR:CD2	2.53	0.42
1:A:485:TYR:HA	1:A:502:ARG:O	2.19	0.42
1:B:492:LEU:HD22	1:B:591:TYR:CD2	2.54	0.42
1:A:507:ILE:HG13	1:A:528:PHE:CE1	2.54	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:537:SER:O	1:A:538:GLU:C	2.64	0.41
1:B:477:THR:HG22	1:B:515:GLU:HB3	2.03	0.41
1:A:478:GLN:HB3	1:A:554:ARG:HB3	2.02	0.41
1:B:483:MET:HG3	1:B:505:TYR:CE1	2.56	0.41
1:A:488:ILE:O	1:A:499:LYS:HB3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	229/239 (96%)	216 (94%)	12 (5%)	1 (0%)	30 60
1	B	233/239 (98%)	211 (91%)	18 (8%)	4 (2%)	7 25
All	All	462/478 (97%)	427 (92%)	30 (6%)	5 (1%)	11 36

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	438	SER
1	B	615	ALA
1	A	425	PRO
1	B	389	THR
1	B	393	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	182/206 (88%)	175 (96%)	7 (4%)	29	64
1	B	181/206 (88%)	174 (96%)	7 (4%)	28	64
All	All	363/412 (88%)	349 (96%)	14 (4%)	28	64

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	389	THR
1	A	391	GLU
1	A	467	CYS
1	A	491	SER
1	A	497	ASP
1	A	506	SER
1	A	584	SER
1	B	410	THR
1	B	413	MET
1	B	467	CYS
1	B	484	GLU
1	B	491	SER
1	B	497	ASP
1	B	506	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	403	HIS
1	B	403	HIS
1	B	455	ASN
1	B	463	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	APR	B	701	-	37,39,39	0.48	0	53,60,60	0.75	1 (1%)
2	APR	A	701	-	37,39,39	0.50	0	53,60,60	0.87	4 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	APR	B	701	-	-	10/22/54/54	0/4/4/4
2	APR	A	701	-	-	4/22/54/54	0/4/4/4

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	701	APR	C1D-C2D-C3D	2.82	105.82	102.30
2	A	701	APR	C1D-C2D-C3D	2.50	105.42	102.30
2	A	701	APR	O2B-PB-O1B	2.30	123.62	112.24
2	A	701	APR	O2D-C2D-C3D	2.24	119.07	111.82
2	A	701	APR	O2A-PA-O1A	2.04	122.34	112.24

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	701	APR	C5D-O5D-PB-O3A
2	A	701	APR	C5D-O5D-PB-O2B
2	B	701	APR	C5D-O5D-PB-O1B
2	B	701	APR	C5D-O5D-PB-O2B
2	B	701	APR	O4D-C4D-C5D-O5D
2	B	701	APR	O4'-C4'-C5'-O5'
2	B	701	APR	C3D-C4D-C5D-O5D
2	B	701	APR	C4D-C5D-O5D-PB
2	B	701	APR	C3'-C4'-C5'-O5'
2	B	701	APR	C5D-O5D-PB-O3A
2	B	701	APR	C2'-C1'-N9-C8
2	A	701	APR	O4'-C4'-C5'-O5'
2	A	701	APR	PB-O3A-PA-O2A
2	B	701	APR	O4'-C1'-N9-C8


There are no ring outliers.

2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	701	APR	1	0
2	A	701	APR	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand APR B 701




Bond lengths



Bond angles




Torsions



Rings

Ligand APR A 701



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	229/239 (95%)	0.05	8 (3%)	47	38	23, 68, 120, 145	2 (0%)
1	B	235/239 (98%)	0.40	17 (7%)	21	16	53, 91, 138, 177	0
All	All	464/478 (97%)	0.23	25 (5%)	31	24	23, 80, 130, 177	2 (0%)

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	617	GLU	5.6
1	B	390	PRO	4.8
1	A	385	LYS	4.6
1	A	384	ALA	4.5
1	B	383	GLN	4.1
1	B	384	ALA	3.9
1	A	611	GLU	3.7
1	B	406	ASP	3.1
1	B	407	GLU	3.1
1	B	389	THR	2.9
1	A	428	THR	2.9
1	B	616	GLN	2.7
1	A	383	GLN	2.7
1	B	403	HIS	2.7
1	A	486[A]	HIS	2.7
1	B	586	LEU	2.6
1	B	425	PRO	2.5
1	A	433	LEU	2.3
1	B	401	VAL	2.3
1	B	426	GLN	2.3
1	B	410	THR	2.3
1	B	398	LEU	2.2
1	B	400	LYS	2.1
1	B	402	ARG	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	427	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	APR	B	701	36/36	0.89	0.11	83,100,117,122	0
2	APR	A	701	36/36	0.95	0.09	53,67,96,112	0
3	ZN	B	703	1/1	0.97	0.04	67,67,67,67	0
3	ZN	A	702	1/1	0.98	0.04	99,99,99,99	0
3	ZN	B	702	1/1	0.99	0.03	72,72,72,72	0
3	ZN	A	703	1/1	0.99	0.03	54,54,54,54	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around APR B 701:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)








Electron density around APR A 701:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers

There are no such residues in this entry.