Full wwPDB X-ray Structure Validation Report (i)

Oct 29, 2026 – 02:05 PM EDT

PDB ID	:	11LZ / pdb_000011lz
Title	:	Crystal structure of Interferon alpha2 bound to IFA001 Fab
Authors	:	Vasquez, S.; Rubio, A.A.; Barnes, C.O.
Deposited on	:	2026-03-04
Resolution	:	2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.2
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	194	3% 63% • 35%
1	B	194	5% 63% 5% 32%
2	C	216	14% 84% 12% •
2	L	216	3% 89% 9% •
3	D	230	10% 79% 13% • 7%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	H	230	

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 8531 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interferon alpha-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	127	Total	C	N	O	S	0	0	0
			1051	679	176	189	7			
1	B	131	Total	C	N	O	S	0	0	0
			1069	686	182	194	7			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	166	GLU	-	expression tag	UNP P01563
A	167	ASN	-	expression tag	UNP P01563
A	168	LEU	-	expression tag	UNP P01563
A	169	TYR	-	expression tag	UNP P01563
A	170	PHE	-	expression tag	UNP P01563
A	171	GLN	-	expression tag	UNP P01563
B	166	GLU	-	expression tag	UNP P01563
B	167	ASN	-	expression tag	UNP P01563
B	168	LEU	-	expression tag	UNP P01563
B	169	TYR	-	expression tag	UNP P01563
B	170	PHE	-	expression tag	UNP P01563
B	171	GLN	-	expression tag	UNP P01563


- Molecule 2 is a protein called IFA001 Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	207	Total	C	N	O	S	0	0	0
			1542	968	262	307	5			
2	L	212	Total	C	N	O	S	0	1	0
			1579	991	268	315	5			

- Molecule 3 is a protein called IFA001 Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	214	Total	C	N	O	S	0	0	0
			1614	1026	269	313	6			
3	H	221	Total	C	N	O	S	0	0	0
			1648	1045	272	325	6			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	D	1	Total	C	N	O	0	0
			14	8	1	5		
4	H	1	Total	C	N	O	0	0
			14	8	1	5		

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Interferon alpha-2

Chain A:




• Molecule 1: Interferon alpha-2

Chain B:




• Molecule 2: IFA001 Fab light chain


Chain C:




• Molecule 2: IFA001 Fab light chain


Chain L:







• Molecule 3: IFA001 Fab heavy chain





• Molecule 3: IFA001 Fab heavy chain





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α, β, γ	37.73Å 63.44Å 130.26Å 100.20° 97.26° 91.06°	Depositor
Resolution (Å)	38.43 – 2.80 38.43 – 2.80	Depositor EDS
% Data completeness (in resolution range)	91.4 (38.43-2.80) 91.3 (38.43-2.80)	Depositor EDS
R_{merge}	0.28	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.18 (at 2.81Å)	Xtriage
Refinement program	PHENIX 1.21.2_5419	Depositor
R, R_{free}	0.221 , 0.279 0.221 , 0.278	Depositor DCC
R_{free} test set	1998 reflections (6.88%)	wwPDB-VP
Wilson B-factor (Å^2)	25.6	Xtriage
Anisotropy	1.497	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.30 , 38.2	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.028 for h,-k,-h-l	Xtriage
F_o,F_c correlation	0.89	EDS
Total number of atoms	8531	wwPDB-VP
Average B, all atoms (Å^2)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.23% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	A	0.19	0/1071	0.30	0/1439
1	B	0.16	0/1088	0.30	0/1462
2	C	0.15	0/1580	0.32	0/2157
2	L	0.21	0/1619	0.38	0/2211
3	D	0.16	0/1656	0.34	0/2257
3	H	0.19	0/1690	0.37	0/2306
All	All	0.18	0/8704	0.34	0/11832

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1051	0	1047	4	0
1	B	1069	0	1063	5	0
2	C	1542	0	1495	14	0
2	L	1579	0	1535	10	0
3	D	1614	0	1545	18	0
3	H	1648	0	1566	5	0
4	D	14	0	13	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	H	14	0	13	0	0
All	All	8531	0	8277	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:35:TRP:HB2	2:C:48:ILE:HB	1.75	0.69
2:L:80:SER:HA	2:L:106:VAL:HG21	1.75	0.69
1:A:117:LEU:HD23	2:L:30:VAL:HG21	1.77	0.65
2:C:43:ALA:HB2	3:D:105:GLN:HA	1.81	0.63
2:C:4:LEU:HD11	2:C:90:THR:HG22	1.79	0.63
3:D:123:PRO:HB3	3:D:211:VAL:HG22	1.82	0.61
2:L:46:LEU:HD21	2:L:49:SER:HB3	1.83	0.61
3:H:82:MET:HB3	3:H:82(C):LEU:HD21	1.84	0.58
3:H:87:THR:HG23	3:H:110:THR:HA	1.85	0.57
3:D:82:MET:HB3	3:D:82(C):LEU:HD21	1.87	0.56
2:L:92:ASP:HB3	2:L:95:LEU:HB2	1.88	0.55
2:C:39:LEU:HD23	2:C:84:ALA:HB2	1.87	0.55
3:D:87:THR:HG23	3:D:110:THR:HA	1.90	0.54
3:D:37:VAL:HG13	3:D:47:TRP:HA	1.89	0.53
2:L:78:LEU:HD11	2:L:104:LEU:HD21	1.90	0.53
2:C:149:LYS:HB2	2:C:192:SER:HB2	1.90	0.53
2:C:27(B):ASN:O	2:C:31:ASN:HB2	2.12	0.49
3:D:154:TRP:HB3	3:D:159:LEU:HD12	1.94	0.49
2:L:140:TYR:HB2	2:L:171:LYS:HG2	1.95	0.49
2:C:125:LEU:HD22	2:C:182:PRO:HG3	1.95	0.49
3:D:83:ARG:HD2	3:D:85:GLU:OE2	2.13	0.48
2:C:32:HIS:HB3	2:C:50:ALA:CB	2.43	0.48
3:D:195:ILE:HG12	3:D:210:ARG:HG3	1.96	0.48
3:H:20:LEU:HD12	3:H:80:LEU:HD23	1.93	0.48
1:A:9:LEU:HD22	1:A:12:ARG:HE	1.79	0.48
3:D:116:THR:HG22	3:D:147:PRO:HD3	1.95	0.47
3:D:17:SER:HA	3:D:82:MET:O	2.14	0.47
3:D:163:VAL:HG22	3:D:182:VAL:HB	1.98	0.46
3:D:12:VAL:HG13	3:D:111:VAL:HG22	1.96	0.46
1:B:59:MET:HE3	1:B:63:ILE:HD11	1.97	0.46
2:C:32:HIS:HB3	2:C:50:ALA:HB1	1.98	0.46
3:D:59:TYR:CZ	3:D:69:ILE:HG22	2.51	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:182:VAL:HG22	3:D:184:VAL:HG13	1.97	0.45
2:C:204:LYS:HD3	2:C:204:LYS:HA	1.83	0.45
3:H:170:LEU:HD13	3:H:176:TYR:CZ	2.52	0.44
3:D:82(C):LEU:HB3	3:D:111:VAL:HG21	1.99	0.44
1:A:12:ARG:HH22	1:B:90:GLN:HG3	1.82	0.44
2:C:46:LEU:HD21	2:C:49:SER:HB2	1.98	0.44
2:C:161:THR:HG23	2:C:176:SER:HB2	1.99	0.44
2:C:194:GLN:HG2	2:C:203:GLU:HG3	2.00	0.44
1:A:60:ILE:HG23	1:A:88:LEU:HD22	2.00	0.44
2:L:204:LYS:HD3	2:L:204:LYS:HA	1.86	0.44
2:L:35:TRP:HB2	2:L:48:ILE:HB	2.00	0.43
2:L:91:TRP:CZ2	2:L:95(A):ASN:HA	2.53	0.42
1:B:14:THR:HG23	1:B:84:PHE:CE1	2.55	0.42
3:D:142:VAL:HG11	3:D:150:VAL:HG11	2.01	0.42
3:D:48:VAL:HG13	3:D:63:VAL:HG21	2.02	0.42
3:D:162:GLY:O	3:D:182:VAL:HA	2.20	0.42
2:C:15:PRO:HA	2:C:78:LEU:HB3	2.01	0.41
1:B:30:LEU:HD22	1:B:33:ARG:HD2	2.01	0.41
3:H:119:PRO:HD2	3:H:205:THR:HG21	2.01	0.41
2:L:46:LEU:HG	2:L:55:PRO:HD2	2.03	0.41
1:B:92:LEU:HD23	1:B:92:LEU:HA	1.93	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	121/194 (62%)	119 (98%)	2 (2%)	0	100	100
1	B	125/194 (64%)	121 (97%)	4 (3%)	0	100	100
2	C	199/216 (92%)	188 (94%)	11 (6%)	0	100	100
2	L	209/216 (97%)	196 (94%)	13 (6%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	D	208/230 (90%)	202 (97%)	6 (3%)	0	100	100
3	H	215/230 (94%)	208 (97%)	7 (3%)	0	100	100
All	All	1077/1280 (84%)	1034 (96%)	43 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	115/177 (65%)	115 (100%)	0	100	100
1	B	117/177 (66%)	116 (99%)	1 (1%)	70	89
2	C	173/182 (95%)	172 (99%)	1 (1%)	78	92
2	L	176/182 (97%)	174 (99%)	2 (1%)	65	87
3	D	176/192 (92%)	172 (98%)	4 (2%)	44	78
3	H	180/192 (94%)	176 (98%)	4 (2%)	45	78
All	All	937/1102 (85%)	925 (99%)	12 (1%)	61	86

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	99	VAL
2	C	167	GLN
3	D	12	VAL
3	D	78	VAL
3	D	182	VAL
3	D	209	LYS
3	H	30	GLU
3	H	63	VAL
3	H	140	CYS
3	H	196	CYS
2	L	19	VAL
2	L	81	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	91	GLN
2	C	17	GLN
2	C	108	GLN
2	C	167	GLN
2	C	170	ASN
3	D	31	ASN
3	D	171	GLN
3	H	164	HIS
2	L	17	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	NAG	H	301	3	14,14,15	0.72	0	17,19,21	1.53	2 (11%)
4	NAG	D	301	3	14,14,15	0.71	0	17,19,21	1.84	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	H	301	3	-	2/6/23/26	0/1/1/1
4	NAG	D	301	3	-	3/6/23/26	0/1/1/1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
4	D	301	NAG	C2-N2-C7	5.50	130.27	122.90
4	H	301	NAG	C1-O5-C5	3.94	117.46	112.19
4	H	301	NAG	C2-N2-C7	3.33	127.36	122.90
4	D	301	NAG	C1-C2-N2	2.66	114.62	110.43
4	D	301	NAG	O7-C7-N2	2.24	125.94	121.98

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	301	NAG	C1-C2-N2-C7
4	D	301	NAG	O5-C5-C6-O6
4	D	301	NAG	C4-C5-C6-O6
4	H	301	NAG	C1-C2-N2-C7
4	H	301	NAG	C3-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	127/194 (65%)	0.52	5 (3%) 43 34	8, 27, 57, 77	0
1	B	131/194 (67%)	0.69	10 (7%) 20 14	14, 35, 63, 75	0
2	C	207/216 (95%)	1.00	31 (14%) 5 4	18, 40, 77, 87	0
2	L	212/216 (98%)	0.29	6 (2%) 55 45	8, 20, 47, 59	1 (0%)
3	D	214/230 (93%)	1.07	23 (10%) 11 8	15, 43, 61, 81	0
3	H	221/230 (96%)	0.36	4 (1%) 67 58	8, 23, 40, 67	0
All	All	1112/1280 (86%)	0.66	79 (7%) 22 16	8, 30, 63, 87	1 (0%)

All (79) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	D	17	SER	4.6
2	C	151	ASP	4.6
3	H	133	GLY	3.8
3	D	159	LEU	3.7
2	C	51	ASN	3.7
2	C	127	ALA	3.5
2	C	120	PRO	3.4
2	L	2	SER	3.4
2	L	55	PRO	3.4
3	D	43	ARG	3.3
1	A	9	LEU	3.3
2	C	189	ARG	3.3
2	C	209	THR	3.3
2	L	52	HIS	3.2
2	C	122	SER	3.0
3	D	82	MET	3.0
2	C	60	ASP	2.9
1	A	43	PHE	2.9
3	D	172	SER	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	D	183	THR	2.9
2	C	179	SER	2.8
3	D	173	SER	2.8
3	H	131	THR	2.8
2	C	128	ASN	2.8
2	C	121	SER	2.8
1	A	41	GLU	2.8
1	B	13	ARG	2.8
2	C	160	GLU	2.8
1	B	41	GLU	2.7
3	H	55	GLU	2.7
2	C	132	LEU	2.7
3	D	138	LEU	2.7
3	D	114	ALA	2.7
2	C	207	ALA	2.6
3	D	156	SER	2.6
2	C	180	LEU	2.6
3	D	212	GLU	2.6
1	B	30	LEU	2.5
1	B	40	GLN	2.5
3	D	144	ASP	2.5
1	A	52	THR	2.5
2	C	116	THR	2.5
3	D	185	PRO	2.4
3	D	186	SER	2.4
3	D	67	PHE	2.4
2	C	42	MET	2.4
3	D	134	GLY	2.4
2	C	131	THR	2.4
2	C	57	GLY	2.4
2	C	67	SER	2.4
1	B	156	ASN	2.3
1	B	43	PHE	2.3
3	D	23	SER	2.3
1	B	26	LEU	2.3
2	C	184	GLN	2.3
1	B	25	SER	2.3
2	C	181	THR	2.3
2	C	123	GLU	2.2
2	C	125	LEU	2.2
2	C	153	SER	2.2
3	D	194	TYR	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	C	182	PRO	2.2
3	H	191	THR	2.2
3	D	184	VAL	2.2
2	L	1	GLN	2.2
2	L	3	VAL	2.2
2	C	124	GLU	2.2
3	D	76	ASN	2.1
2	C	156	LYS	2.1
3	D	213	PRO	2.1
1	B	99	VAL	2.1
1	B	24	ILE	2.1
1	A	29	CYS	2.0
3	D	174	GLY	2.0
2	C	183	GLU	2.0
3	D	142	VAL	2.0
2	C	150	ALA	2.0
2	C	185	TRP	2.0
2	L	30	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	NAG	D	301	14/15	0.58	0.18	56,76,87,88	0
4	NAG	H	301	14/15	0.59	0.24	56,70,79,80	0

6.5 Other polymers (i)

There are no such residues in this entry.