Full wwPDB X-ray Structure Validation Report (i)

Nov 12, 2026 – 02:11 PM JST

PDB ID	:	24YI / pdb_000024yi
Title	:	3-methyladenine DNA glycosylase bound to DNA/THF
Authors	:	Xie, W.; Jia, Q.; Jiang, P.
Deposited on	:	2026-03-24
Resolution	:	1.94 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.2
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1452 (1.94-1.94)
Clashscore	190562	1494 (1.94-1.94)
Ramachandran outliers	187476	1479 (1.94-1.94)
Sidechain outliers	187428	1479 (1.94-1.94)
RSRZ outliers	180081	1453 (1.94-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	286	3%, 91%, 5%
2	C	16	88%, 12%
3	I	16	56%, 31%, 12%

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 2981 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Base excision DNA repair protein, HhH-GPD superfamily.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	275	Total	C	N	O	S	0	0	0
			2194	1422	362	405	5			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	279	LEU	-	expression tag	UNP C5A632
A	280	GLU	-	expression tag	UNP C5A632
A	281	HIS	-	expression tag	UNP C5A632
A	282	HIS	-	expression tag	UNP C5A632
A	283	HIS	-	expression tag	UNP C5A632
A	284	HIS	-	expression tag	UNP C5A632
A	285	HIS	-	expression tag	UNP C5A632
A	286	HIS	-	expression tag	UNP C5A632

- Molecule 2 is a DNA chain called DNA (5'-D(*TP*CP*GP*CP*TP*AP*CP*AP*TP*GP*TP*CP*GP*TP*CP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	16	Total	C	N	O	P	0	0	0
			321	155	55	96	15			

- Molecule 3 is a DNA chain called DNA (5'-D(*TP*GP*AP*CP*GP*AP*CP*(3DR)P*TP*GP*TP*AP*GP*CP*GP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	I	16	Total	C	N	O	P	0	0	0
			319	152	60	92	15			

- Molecule 4 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Na 1 1	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	116	Total O 116 116	0	0
5	C	9	Total O 9 9	0	0
5	I	21	Total O 21 21	0	0


3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Base excision DNA repair protein, HhH-GPD superfamily

Chain A:





• Molecule 2: DNA (5'-D(*TP*CP*GP*CP*TP*AP*CP*AP*TP*GP*TP*CP*GP*TP*CP*A)-3')

Chain C:






• Molecule 3: DNA (5'-D(*TP*GP*AP*CP*GP*AP*CP*(3DR)P*TP*GP*TP*AP*GP*CP*GP*A)-3')

Chain I:





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	45.10Å 88.54Å 91.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.67 – 1.94 40.67 – 1.94	Depositor EDS
% Data completeness (in resolution range)	94.3 (40.67-1.94) 94.3 (40.67-1.94)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.59 (at 1.94Å)	Xtriage
Refinement program	PHENIX 1.14_3260	Depositor
R, R_{free}	0.176 , 0.214 0.177 , 0.216	Depositor DCC
R_{free} test set	1282 reflections (4.57%)	wwPDB-VP
Wilson B-factor (Å^2)	35.4	Xtriage
Anisotropy	0.312	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.35 , 34.0	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	0.017 for -h,l,k	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	2981	wwPDB-VP
Average B, all atoms (Å^2)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.07% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 3DR, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|$ >5	RMSZ	#$\|Z\|$ >5
1	A	0.45	0/2247	0.65	2/3039 (0.1%)
2	C	1.08	0/358	1.07	0/550
3	I	1.26	2/345 (0.6%)	1.13	1/529 (0.2%)
All	All	0.69	2/2950 (0.1%)	0.80	3/4118 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	I	16	DA	C6-N6	6.53	1.39	1.33
3	I	9	DT	C5'-C4'	5.33	1.55	1.51

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	122	ARG	CG-CD-NE	-11.86	85.90	112.00
3	I	14	DC	O4'-C1'-N1	5.35	111.75	108.00
1	A	122	ARG	N-CA-CB	-5.19	102.49	110.01

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	48	GLU	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2194	0	2140	17	1
2	C	321	0	183	1	0
3	I	319	0	178	9	1
4	A	1	0	0	0	0
5	A	116	0	0	2	0
5	C	9	0	0	0	0
5	I	21	0	0	1	0
All	All	2981	0	2501	27	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:I:8:3DR:O4'	3:I:8:3DR:C4'	1.65	1.18
1:A:118:ASP:O	1:A:122:ARG:HG2	1.80	0.81
2:C:14:DT:H2"	2:C:15:DC:H5"	1.68	0.75
1:A:122:ARG:HH11	1:A:122:ARG:HA	1.55	0.71
3:I:8:3DR:OP1	5:I:101:HOH:O	2.12	0.68
1:A:122:ARG:HB3	1:A:122:ARG:CZ	2.25	0.67
1:A:119:PHE:O	1:A:122:ARG:HB2	1.95	0.66
1:A:122:ARG:HB3	1:A:122:ARG:NH1	2.13	0.63
1:A:122:ARG:HH11	1:A:122:ARG:CA	2.12	0.62
3:I:8:3DR:O4'	3:I:8:3DR:C5'	2.45	0.62
1:A:40:TYR:OH	5:A:401:HOH:O	2.17	0.57
1:A:122:ARG:CZ	1:A:122:ARG:CB	2.84	0.56
1:A:9:LYS:HE3	1:A:270:LYS:NZ	2.21	0.55
1:A:122:ARG:NH1	1:A:122:ARG:CB	2.72	0.51
3:I:15:DG:C6	3:I:16:DA:C6	2.99	0.50
3:I:4:DC:H2"	3:I:5:DG:C8	2.47	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:I:15:DG:C4	3:I:16:DA:N7	2.80	0.49
3:I:15:DG:C4	3:I:16:DA:C8	3.00	0.49
1:A:96:MET:HE1	5:A:503:HOH:O	2.13	0.48
3:I:15:DG:H2"	3:I:16:DA:H8	1.77	0.48
1:A:270:LYS:O	1:A:274:GLU:HG2	2.15	0.46
1:A:122:ARG:NH1	1:A:122:ARG:CA	2.78	0.46
3:I:8:3DR:C5'	3:I:8:3DR:C1'	2.81	0.46
1:A:122:ARG:HA	1:A:122:ARG:HD2	1.70	0.45
1:A:30:LEU:HD21	1:A:61:LYS:HD2	1.99	0.44
1:A:187:ASP:OD1	1:A:190:GLU:HG3	2.20	0.42
1:A:36:ARG:HB2	1:A:50:TRP:CZ2	2.54	0.42

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:122:ARG:NH2	3:I:16:DA:N7[1_655]	2.17	0.03

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	273/286 (96%)	265 (97%)	6 (2%)	2 (1%)	18 9

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	49	ASP
1	A	48	GLU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	218/238 (92%)	218 (100%)	0	100 100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	45	HIS
1	A	104	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	3DR	I	8	3	8,11,12	5.56	6 (75%)	9,14,17	1.26	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	3DR	I	8	3	-	2/3/15/16	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	I	8	3DR	O4'-C4'	12.65	1.65	1.44
3	I	8	3DR	C3'-C4'	-6.63	1.34	1.53
3	I	8	3DR	O4'-C1'	-3.89	1.31	1.42
3	I	8	3DR	C2'-C1'	3.26	1.60	1.51
3	I	8	3DR	O5'-C5'	-2.66	1.38	1.44
3	I	8	3DR	C2'-C3'	2.27	1.56	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	I	8	3DR	C1'-O4'-C4'	-2.96	103.71	108.48

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	I	8	3DR	C3'-C4'-C5'-O5'
3	I	8	3DR	O4'-C4'-C5'-O5'

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	I	8	3DR	4	0

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	275/286 (96%)	0.12	8 (2%) 53 60	25, 35, 62, 83	0
2	C	16/16 (100%)	-0.01	0 100 100	41, 51, 69, 69	0
3	I	15/16 (93%)	0.02	0 100 100	33, 53, 60, 62	0
All	All	306/318 (96%)	0.11	8 (2%) 57 64	25, 36, 65, 83	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	122	ARG	3.7
1	A	31	GLY	3.2
1	A	48	GLU	2.9
1	A	275	ARG	2.7
1	A	49	ASP	2.6
1	A	32	GLY	2.6
1	A	50	TRP	2.5
1	A	46	PHE	2.2

6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	3DR	I	8	11/12	0.95	0.08	29,40,48,51	0

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	NA	A	301	1/1	0.98	0.03	28,28,28,28	0

6.5 Other polymers (i)

There are no such residues in this entry.