



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 19, 2026 – 12:36 PM EDT

PDB ID : 36AC / pdb_000036ac
Title : Ternary complex between CHD4-C1bC2ab, GATAD2ACR2, and CDK2AP1
Authors : Williams, D.C.; Li, T.; McGinty, R.K.
Deposited on : 2026-05-27
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.2
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	89	
2	A	340	
3	B	154	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4153 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cyclin-dependent kinase 2-associated protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	66	Total	C	N	O	S	0	0	0
			481	305	87	87	2			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	27	GLY	-	expression tag	UNP O14519

- Molecule 2 is a protein called Chromodomain-helicase-DNA-binding protein 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	327	Total	C	N	O	S	0	0	0
			2512	1598	460	441	13			

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1394	MET	-	initiating methionine	UNP B3KY63
A	1520	GLY	-	linker	UNP B3KY63
A	1521	GLY	-	linker	UNP B3KY63
A	1522	GLY	-	linker	UNP B3KY63
A	1523	SER	-	linker	UNP B3KY63

- Molecule 3 is a protein called Transcriptional repressor p66-alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	B	131	Total	C	N	O	S	0	0	0
			920	583	156	172	9			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	333	GLY	-	expression tag	UNP Q86YP4
B	334	SER	-	expression tag	UNP Q86YP4

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total Zn 1 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	C	32	Total O 32 32	0	0
5	A	170	Total O 170 170	0	0
5	B	37	Total O 37 37	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cyclin-dependent kinase 2-associated protein 1



- Molecule 2: Chromodomain-helicase-DNA-binding protein 4



- Molecule 3: Transcriptional repressor p66-alpha



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	43.73Å 52.86Å 72.47Å 98.75° 103.40° 110.69°	Depositor
Resolution (Å)	30.03 – 2.00 30.03 – 2.00	Depositor EDS
% Data completeness (in resolution range)	94.3 (30.03-2.00) 94.3 (30.03-2.00)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.11 (at 2.01Å)	Xtriage
Refinement program	PHENIX 2.0_5885	Depositor
R, R_{free}	0.197 , 0.245 0.198 , 0.246	Depositor DCC
R_{free} test set	2000 reflections (5.22%)	wwPDB-VP
Wilson B-factor (Å ²)	39.0	Xtriage
Anisotropy	0.334	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 61.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4153	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	C	0.26	0/487	0.46	0/655
2	A	0.34	0/2570	0.48	0/3481
3	B	0.27	0/935	0.43	0/1275
All	All	0.32	0/3992	0.47	0/5411

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	481	0	465	4	0
2	A	2512	0	2397	21	0
3	B	920	0	819	7	0
4	B	1	0	0	0	0
5	A	170	0	0	1	0
5	B	37	0	0	1	0
5	C	32	0	0	0	0
All	All	4153	0	3681	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:1472:ALA:HB1	2:A:1475:ALA:HB3	1.80	0.63
1:C:95:ILE:HD11	2:A:1864:MET:HE2	1.90	0.54
1:C:45:PRO:HD2	2:A:1765:PHE:HE2	1.74	0.53
1:C:112:ASN:HB2	2:A:1847:ASN:HD22	1.74	0.52
2:A:1493:LEU:HD12	3:B:386:TYR:CE1	2.44	0.52
2:A:1714:ASN:HD22	2:A:1717:ARG:HH12	1.58	0.51
2:A:1813:HIS:CE1	2:A:1878:ILE:HD11	2.46	0.51
2:A:1725:THR:O	2:A:1730:HIS:HE1	1.95	0.49
2:A:1696:PHE:CZ	2:A:1732:ARG:HA	2.48	0.49
2:A:1821:ARG:HD3	5:A:2094:HOH:O	2.16	0.46
2:A:1813:HIS:NE2	2:A:1878:ILE:HD11	2.31	0.46
2:A:1845:PRO:HB2	3:B:356:LEU:HD12	1.98	0.46
2:A:1502:ILE:HG22	2:A:1742:ILE:HG12	1.99	0.43
3:B:373:MET:SD	3:B:376:LEU:HD23	2.58	0.43
2:A:1821:ARG:HB3	2:A:1864:MET:HG2	2.00	0.43
2:A:1768:GLU:HG2	2:A:1774:PHE:CG	2.54	0.42
2:A:1851:HIS:HA	2:A:1854:LEU:HD12	2.02	0.41
2:A:1742:ILE:HD13	2:A:1742:ILE:HA	1.80	0.41
2:A:1433:MET:HE2	3:B:396:ASN:CG	2.45	0.41
1:C:112:ASN:HB2	2:A:1847:ASN:ND2	2.36	0.41
3:B:429:ARG:HG3	5:B:619:HOH:O	2.20	0.41
2:A:1493:LEU:HD12	3:B:386:TYR:HE1	1.87	0.40
2:A:1704:GLY:N	3:B:426:CYS:HB2	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	62/89 (70%)	60 (97%)	2 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	323/340 (95%)	318 (98%)	5 (2%)	0	100	100
3	B	127/154 (82%)	125 (98%)	2 (2%)	0	100	100
All	All	512/583 (88%)	503 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	43/72 (60%)	43 (100%)	0	100	100
2	A	238/288 (83%)	237 (100%)	1 (0%)	84	89
3	B	80/129 (62%)	80 (100%)	0	100	100
All	All	361/489 (74%)	360 (100%)	1 (0%)	86	91

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	1742	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	37	GLN
2	A	1714	ASN
2	A	1753	GLN
2	A	1762	ASN
2	A	1773	ASN
2	A	1834	HIS
3	B	419	GLN
3	B	440	ASN
3	B	445	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	C	66/89 (74%)	0.61	1 (1%) 72 71	34, 53, 85, 102	0
2	A	327/340 (96%)	0.35	15 (4%) 37 36	27, 47, 78, 100	0
3	B	131/154 (85%)	0.51	9 (6%) 23 21	33, 60, 91, 104	0
All	All	524/583 (89%)	0.42	25 (4%) 35 34	27, 51, 85, 104	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	A	1479	ALA	4.9
1	C	34	GLN	3.3
3	B	343	SER	3.2
3	B	345	GLN	2.9
2	A	1767	GLY	2.7
2	A	1842	GLY	2.7
3	B	371	PRO	2.7
3	B	370	ALA	2.6
3	B	347	ALA	2.6
2	A	1472	ALA	2.6
2	A	1408	GLY	2.5
2	A	1434	PRO	2.4
2	A	1726	TYR	2.3
2	A	1897	ALA	2.3
2	A	1725	THR	2.3
2	A	1807	MET	2.3
2	A	1721	VAL	2.2
3	B	368	PRO	2.2
2	A	1436	GLN	2.2
3	B	409	VAL	2.2
2	A	1435	PRO	2.1
3	B	400	THR	2.2
2	A	1487	LEU	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	A	1440	THR	2.0
3	B	477	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	ZN	B	501	1/1	0.99	0.03	45,45,45,45	0

6.5 Other polymers [i](#)

There are no such residues in this entry.