Full wwPDB X-ray Structure Validation Report (i)

Oct 26, 2026 – 10:24 AM EDT

PDB ID	:	36OU / pdb_000036ou
Title	:	Room-temperature multiconformer model of PSD-95 PDZ3-CRIPT
Authors	:	Wang, H.K.; Hekstra, D.R.; White, K.I.
Deposited on	:	2026-06-24
Resolution	:	1.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.2
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1216 (1.36-1.36)
Clashscore	190562	1232 (1.36-1.36)
Ramachandran outliers	187476	1220 (1.36-1.36)
Sidechain outliers	187428	1220 (1.36-1.36)
RSRZ outliers	180081	1214 (1.36-1.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	119	4% 82% 13%
2	B	9	56% 44% 33% 22%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 3104 atoms, of which 1488 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Disks large homolog 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
1	A	117	2686	833	1324	248	281	0	57	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	297	GLY	-	expression tag	UNP P31016
A	298	SER	-	expression tag	UNP P31016
A	299	PRO	-	expression tag	UNP P31016
A	300	GLU	-	expression tag	UNP P31016
A	301	PHE	-	expression tag	UNP P31016
A	403	ASN	-	expression tag	UNP P31016
A	404	SER	-	expression tag	UNP P31016
A	405	ARG	-	expression tag	UNP P31016
A	406	VAL	-	expression tag	UNP P31016
A	407	ASN	-	expression tag	UNP P31016
A	408	SER	-	expression tag	UNP P31016
A	409	SER	-	expression tag	UNP P31016
A	410	GLY	-	expression tag	UNP P31016
A	411	ARG	-	expression tag	UNP P31016
A	412	ILE	-	expression tag	UNP P31016
A	413	VAL	-	expression tag	UNP P31016
A	414	THR	-	expression tag	UNP P31016
A	415	ASP	-	expression tag	UNP P31016

- Molecule 2 is a protein called Cysteine-rich PDZ-binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
2	B	9	321	96	164	28	33	0	8	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	85	Total O 86 86	0	1
3	B	11	Total O 11 11	0	1

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Disks large homolog 4



• Molecule 2: Cysteine-rich PDZ-binding protein



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 41 3 2	Depositor
Cell constants a, b, c, α, β, γ	90.73Å 90.73Å 90.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.58 – 1.35 40.58 – 1.35	Depositor EDS
% Data completeness (in resolution range)	90.8 (40.58-1.35) 90.7 (40.58-1.35)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.63 (at 1.35Å)	Xtriage
Refinement program	PHENIX 2.0_5936	Depositor
R, R_{free}	0.122 , 0.147 0.122 , 0.149	Depositor DCC
R_{free} test set	1500 reflections (5.23%)	wwPDB-VP
Wilson B-factor (Å^2)	14.2	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.41 , 50.5	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.98	EDS
Total number of atoms	3104	wwPDB-VP
Average B, all atoms (Å^2)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.67% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	A	0.61	0/1379	0.68	4/1857 (0.2%)
2	B	0.34	0/158	0.64	0/208
All	All	0.59	0/1537	0.68	4/2065 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	334[A]	GLU	CA-C-N	5.27	131.74	121.41
1	A	334[A]	GLU	C-N-CA	5.27	131.74	121.41
1	A	334[B]	GLU	CA-C-N	5.27	131.74	121.41
1	A	334[B]	GLU	C-N-CA	5.27	131.74	121.41

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	331[A]	GLU	Peptide
1	A	332[A]	ASP	Peptide
1	A	334[A]	GLU	Mainchain
1	A	334[B]	GLU	Mainchain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1362	1324	1302	18	0
2	B	157	164	164	10	0
3	A	86	0	0	3	0
3	B	11	0	0	2	0
All	All	1616	1488	1466	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:331[B]:GLU:OE2	2:B:5[B]:LYS:HE3	1.13	1.22
1:A:331[B]:GLU:OE2	2:B:5[B]:LYS:CE	1.94	1.13
1:A:396[A]:GLU:OE1	3:A:502:HOH:O	1.93	0.84
1:A:377[B]:ILE:HG12	1:A:381:ASN:ND2	1.93	0.83
1:A:331[B]:GLU:OE2	2:B:5[B]:LYS:CD	2.39	0.71
1:A:330[A]:GLY:C	1:A:332[A]:ASP:HB3	2.21	0.65
2:B:5[A]:LYS:HG2	3:B:105:HOH:O	1.97	0.63
1:A:326[B]:ASN:HD22	2:B:6[B]:GLN:HG2	1.64	0.63
1:A:331[A]:GLU:N	1:A:332[A]:ASP:HB3	2.14	0.62
1:A:377[B]:ILE:HG12	1:A:381:ASN:HD22	1.70	0.56
1:A:374:GLN:HA	1:A:377[B]:ILE:HG22	1.94	0.50
1:A:391[B]:GLN:OE1	1:A:393[B]:LYS:NZ	2.45	0.49
2:B:2[A]:LYS:NZ	2:B:2[A]:LYS:HA	2.27	0.49
1:A:355[B]:LYS:HE3	3:A:559:HOH:O	2.13	0.48
1:A:338:ILE:HD11	1:A:359[A]:ILE:HD11	1.97	0.46
1:A:404[B]:SER:OG	1:A:414[B]:THR:HG22	2.16	0.45
1:A:326[B]:ASN:ND2	2:B:6[B]:GLN:HG2	2.29	0.45
1:A:409[B]:SER:HB2	1:A:411:ARG:HH11	1.83	0.43
2:B:3[A]:ASN:ND2	3:B:102:HOH:O	2.51	0.43
1:A:332[B]:ASP:N	1:A:332[B]:ASP:OD1	2.51	0.43
2:B:1[B]:THR:C	2:B:3[B]:ASN:N	2.76	0.42
2:B:1[B]:THR:C	2:B:3[B]:ASN:H	2.27	0.42
1:A:414[B]:THR:HG23	3:A:533:HOH:O	2.19	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	175/119 (147%)	163 (93%)	12 (7%)	0	100 100
2	B	16/9 (178%)	14 (88%)	2 (12%)	0	100 100
All	All	191/128 (149%)	177 (93%)	14 (7%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	148/94 (157%)	146 (99%)	2 (1%)	59 28
2	B	19/9 (211%)	15 (79%)	4 (21%)	1 0
All	All	167/103 (162%)	161 (96%)	6 (4%)	37 5

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	399[A]	ARG
1	A	399[B]	ARG
2	B	2[A]	LYS
2	B	2[B]	LYS
2	B	3[A]	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	3[B]	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	374	GLN
1	A	381	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	117/119 (98%)	-0.39	5 (4%) 40 50	5, 14, 28, 66	57 (48%)
2	B	9/9 (100%)	2.95	5 (55%) 0 0	6, 12, 48, 50	8 (88%)
All	All	126/128 (98%)	-0.15	10 (7%) 18 25	5, 14, 32, 66	65 (51%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	3[A]	ASN	8.4
2	B	2[A]	LYS	7.6
2	B	1[A]	THR	5.8
1	A	299	PRO	4.8
2	B	4[A]	TYR	4.3
2	B	5[A]	LYS	3.7
1	A	332[A]	ASP	3.0
1	A	333[A]	GLY	2.7
1	A	304[A]	GLU	2.6
1	A	301[A]	PHE	2.4

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no oligosaccharides in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.