



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 28, 2026 – 12:08 AM UTC

PDB ID : 7QUY / pdb_00007quy
Title : Alcohol Dehydrogenase from Thauera aromatica complexed with NADH
Authors : Petchey, M.L.; Stark, F.; Ansorge-Schumacher, M.; Grogan, G.
Deposited on : 2022-01-19
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION




The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	354	
1	B	354	
1	C	354	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ZN	B	405	-	-	X	-

2 Entry composition ⓘ


There are 5 unique types of molecules in this entry. The entry contains 7807 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 6-hydroxycyclohex-1-ene-1-carbonyl-CoA dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	352	Total	C	N	O	S	0	0	0
			2532	1596	450	468	18			
1	B	352	Total	C	N	O	S	0	0	0
			2541	1603	451	469	18			
1	C	352	Total	C	N	O	S	0	0	0
			2473	1563	434	458	18			

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (CCD ID: NAD) (formula: C₂₁H₂₇N₇O₁₄P₂) (labeled as "Ligand of Interest" by depositor).



- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Zn	0	0
			2	2		
4	B	2	Total	Zn	0	0
			2	2		
4	C	2	Total	Zn	0	0
			2	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	43	Total	O	0	0
			43	43		
5	B	40	Total	O	0	0
			40	40		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	20	Total	O	0	0
			20	20		

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	75.42Å 239.33Å 168.46Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	84.37 – 2.60 84.37 – 2.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (84.37-2.60) 100.0 (84.37-2.60)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.10 (at 2.62Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.215 , 0.252 0.219 , 0.250	Depositor DCC
R_{free} test set	2352 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	58.7	Xtriage
Anisotropy	0.739	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 40.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7807	wwPDB-VP
Average B, all atoms (Å ²)	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, NAD, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.07	0/2583	1.48	7/3527 (0.2%)
1	B	1.05	0/2592	1.44	1/3537 (0.0%)
1	C	1.10	0/2524	1.46	2/3455 (0.1%)
All	All	1.08	0/7699	1.46	10/10519 (0.1%)

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	94	CYS	CB-CA-C	-8.55	96.14	110.68
1	B	94	CYS	CB-CA-C	-7.47	98.86	110.81
1	A	108	CYS	CA-CB-SG	-7.28	97.65	114.40
1	A	246	LEU	N-CA-C	-6.47	104.96	112.92
1	A	107	ILE	CA-C-N	-6.01	112.18	120.71
1	A	107	ILE	C-N-CA	-6.01	112.18	120.71
1	A	100	CYS	CA-CB-SG	6.00	128.19	114.40
1	C	93	PRO	CA-C-N	5.82	128.07	120.28
1	C	93	PRO	C-N-CA	5.82	128.07	120.28
1	A	107	ILE	O-C-N	5.51	129.30	122.14

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2532	0	2473	28	0
1	B	2541	0	2494	33	0
1	C	2473	0	2367	25	0
2	A	44	0	26	3	0
2	B	44	0	26	1	0
2	C	44	0	26	0	0
3	A	10	0	0	0	0
3	B	10	0	0	0	0
4	A	2	0	0	0	0
4	B	2	0	0	4	0
4	C	2	0	0	0	0
5	A	43	0	0	0	0
5	B	40	0	0	1	0
5	C	20	0	0	1	0
All	All	7807	0	7412	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:94:CYS:SG	1:B:96:THR:N	2.20	1.14
1:A:94:CYS:SG	1:A:96:THR:N	2.27	1.07
1:B:94:CYS:SG	1:B:95:GLY:N	2.28	1.07
1:A:94:CYS:SG	1:A:95:GLY:N	2.38	0.96
1:B:108:CYS:HG	4:B:405:ZN:ZN	0.81	0.91
1:B:190:GLN:HE22	1:B:322:ASP:H	1.31	0.78
1:C:190:GLN:HE22	1:C:322:ASP:H	1.31	0.76
1:A:190:GLN:HE22	1:A:322:ASP:H	1.32	0.75
1:B:108:CYS:SG	4:B:405:ZN:ZN	1.76	0.74
1:B:100:CYS:HG	4:B:405:ZN:ZN	0.99	0.74
1:B:153:VAL:HG21	1:B:327:ILE:HD11	1.70	0.72
1:C:153:VAL:HG21	1:C:327:ILE:HD11	1.71	0.72
1:C:94:CYS:SG	1:C:97:CYS:N	2.62	0.71
1:A:80:GLN:HE21	1:A:80:GLN:H	1.40	0.68
1:A:153:VAL:HG21	1:A:327:ILE:HD11	1.77	0.66
1:B:100:CYS:SG	4:B:405:ZN:ZN	1.88	0.63
1:C:239:ALA:HA	1:C:244:LEU:HD12	1.82	0.60
1:C:177:VAL:HG12	1:C:179:ILE:CD1	2.31	0.60
1:C:177:VAL:HG12	1:C:179:ILE:HD11	1.83	0.58
1:C:113:MET:HE2	1:C:116:ASN:HB2	1.86	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:401:NAD:N7N	1:B:292:MET:HE2	2.20	0.57
1:A:190:GLN:NE2	1:A:322:ASP:H	2.03	0.56
1:B:190:GLN:NE2	1:B:322:ASP:H	2.03	0.54
1:A:227:SER:O	1:A:231:LEU:HB2	2.07	0.54
1:B:227:SER:O	1:B:231:LEU:HB2	2.07	0.54
2:A:401:NAD:H2N	1:B:292:MET:CE	2.39	0.53
1:C:179:ILE:HD12	1:C:179:ILE:N	2.22	0.53
1:C:227:SER:O	1:C:231:LEU:HB2	2.08	0.53
1:C:190:GLN:NE2	1:C:322:ASP:H	2.03	0.52
1:A:94:CYS:SG	1:A:97:CYS:N	2.73	0.52
1:B:289:SER:O	1:B:292:MET:HG3	2.10	0.52
1:A:289:SER:O	1:A:292:MET:HG3	2.10	0.52
1:A:292:MET:HE2	2:B:401:NAD:N7N	2.26	0.51
1:B:138:GLU:HB2	1:C:142:ALA:HB1	1.94	0.49
1:B:163:GLN:HG2	1:B:301:ASN:OD1	2.13	0.49
1:A:287:ARG:NH1	1:B:281:MET:O	2.45	0.48
1:C:62:ALA:H	1:C:119:GLN:HE22	1.62	0.48
1:B:100:CYS:HG	1:B:108:CYS:HG	1.60	0.48
1:C:48:TYR:OH	1:C:113:MET:CE	2.61	0.48
1:B:94:CYS:SG	1:B:95:GLY:C	2.93	0.48
1:B:94:CYS:SG	1:B:97:CYS:N	2.75	0.48
1:B:204:VAL:HB	1:B:224:ARG:CZ	2.44	0.48
1:C:48:TYR:OH	1:C:113:MET:HE2	2.14	0.47
1:A:94:CYS:SG	1:A:100:CYS:HB2	2.55	0.47
1:A:160:THR:HG21	2:A:401:NAD:C4N	2.45	0.47
1:A:295:HIS:CD2	1:B:302:TRP:H	2.33	0.47
1:C:245:ARG:HG2	1:C:248:ARG:NH1	2.29	0.47
1:C:163:GLN:HG2	1:C:301:ASN:OD1	2.15	0.46
1:B:240:LYS:NZ	5:B:501:HOH:O	2.48	0.46
1:C:62:ALA:H	1:C:119:GLN:NE2	2.14	0.45
1:A:246:LEU:HD13	1:A:246:LEU:HA	1.82	0.45
1:A:163:GLN:HG2	1:A:301:ASN:OD1	2.17	0.45
1:A:302:TRP:H	1:B:295:HIS:CD2	2.35	0.45
1:B:94:CYS:HB3	1:B:108:CYS:SG	2.57	0.45
1:A:62:ALA:H	1:A:119:GLN:NE2	2.15	0.45
1:B:62:ALA:H	1:B:119:GLN:NE2	2.15	0.44
1:A:94:CYS:SG	1:A:95:GLY:C	2.99	0.43
1:A:153:VAL:O	1:A:157:ALA:HB3	2.18	0.43
1:A:300:GLY:HA3	1:B:291:LEU:O	2.18	0.43
1:B:94:CYS:SG	1:B:100:CYS:CB	3.07	0.43
1:B:170:VAL:HG22	1:B:250:LYS:HG3	1.99	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:62:ALA:H	1:A:119:GLN:HE22	1.65	0.43
1:B:98:GLU:HA	1:B:101:THR:HG22	2.01	0.43
1:B:39:CYS:HA	1:B:66:GLU:O	2.19	0.42
1:A:42:CYS:HB2	1:A:66:GLU:OE2	2.19	0.42
1:A:170:VAL:HG22	1:A:250:LYS:HG3	2.00	0.42
1:C:39:CYS:HA	1:C:66:GLU:O	2.19	0.42
1:A:39:CYS:HA	1:A:66:GLU:O	2.20	0.41
1:B:62:ALA:H	1:B:119:GLN:HE22	1.67	0.41
1:C:178:VAL:C	1:C:179:ILE:HD12	2.45	0.41
1:A:190:GLN:HB3	1:A:321:ILE:HG22	2.02	0.41
1:C:126:VAL:HA	5:C:507:HOH:O	2.21	0.41
1:C:170:VAL:HG22	1:C:250:LYS:HG3	2.03	0.41
1:C:171:GLU:O	1:C:174:ASP:HB2	2.21	0.41
1:C:190:GLN:HB3	1:C:321:ILE:HG22	2.02	0.41
1:B:291:LEU:HD12	1:B:291:LEU:HA	1.88	0.41
1:C:59:LEU:H	1:C:59:LEU:HD22	1.86	0.41
1:C:113:MET:HE2	1:C:116:ASN:CB	2.50	0.41
1:A:141:LEU:HD11	1:A:148:LEU:HA	2.03	0.41
1:B:94:CYS:SG	1:B:95:GLY:CA	3.07	0.40
1:A:179:ILE:CD1	1:A:264:ALA:HB2	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	350/354 (99%)	341 (97%)	8 (2%)	1 (0%)	36	58
1	B	350/354 (99%)	341 (97%)	8 (2%)	1 (0%)	36	58
1	C	350/354 (99%)	339 (97%)	10 (3%)	1 (0%)	36	58
All	All	1050/1062 (99%)	1021 (97%)	26 (2%)	3 (0%)	36	58

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	116	ASN
1	C	116	ASN
1	A	116	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	248/264 (94%)	227 (92%)	21 (8%)	10	22
1	B	250/264 (95%)	231 (92%)	19 (8%)	12	27
1	C	234/264 (89%)	212 (91%)	22 (9%)	8	18
All	All	732/792 (92%)	670 (92%)	62 (8%)	10	22

All (62) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	ARG
1	A	34	VAL
1	A	59	LEU
1	A	80	GLN
1	A	101	THR
1	A	108	CYS
1	A	109	ARG
1	A	112	VAL
1	A	141	LEU
1	A	146	LEU
1	A	178	VAL
1	A	208	LYS
1	A	219	LEU
1	A	245	ARG
1	A	246	LEU
1	A	263	SER
1	A	288	LEU
1	A	291	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	327	ILE
1	A	334	GLN
1	A	349	ARG
1	B	7	ARG
1	B	34	VAL
1	B	59	LEU
1	B	101	THR
1	B	110	ASP
1	B	112	VAL
1	B	141	LEU
1	B	146	LEU
1	B	178	VAL
1	B	219	LEU
1	B	240	LYS
1	B	245	ARG
1	B	246	LEU
1	B	288	LEU
1	B	289	SER
1	B	291	LEU
1	B	319	LYS
1	B	327	ILE
1	B	334	GLN
1	C	7	ARG
1	C	9	MET
1	C	18	VAL
1	C	34	VAL
1	C	59	LEU
1	C	80	GLN
1	C	82	LEU
1	C	84	ARG
1	C	101	THR
1	C	131	ARG
1	C	141	LEU
1	C	146	LEU
1	C	178	VAL
1	C	219	LEU
1	C	227	SER
1	C	245	ARG
1	C	246	LEU
1	C	263	SER
1	C	288	LEU
1	C	291	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	327	ILE
1	C	334	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	56	ASN
1	A	80	GLN
1	A	119	GLN
1	A	167	GLN
1	A	190	GLN
1	A	261	GLN
1	A	330	HIS
1	B	6	HIS
1	B	56	ASN
1	B	119	GLN
1	B	167	GLN
1	B	190	GLN
1	B	330	HIS
1	C	56	ASN
1	C	111	GLN
1	C	119	GLN
1	C	167	GLN
1	C	190	GLN
1	C	330	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 13 ligands modelled in this entry, 6 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAD	A	401	-	46,48,48	0.58	1 (2%)	64,73,73	0.81	1 (1%)
3	SO4	A	403	-	4,4,4	0.27	0	6,6,6	0.09	0
3	SO4	B	402	-	4,4,4	0.35	0	6,6,6	0.10	0
3	SO4	A	402	-	4,4,4	0.46	0	6,6,6	0.08	0
3	SO4	B	403	-	4,4,4	0.28	0	6,6,6	0.08	0
2	NAD	C	401	-	46,48,48	0.51	0	64,73,73	0.80	1 (1%)
2	NAD	B	401	-	46,48,48	0.60	0	64,73,73	0.77	1 (1%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	A	401	-	-	2/30/62/62	0/5/5/5
2	NAD	B	401	-	-	7/30/62/62	0/5/5/5
2	NAD	C	401	-	-	11/30/62/62	0/5/5/5

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	NAD	C2N-N1N	2.05	1.37	1.35

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	NAD	C6N-N1N-C2N	-3.56	118.85	121.88
2	C	401	NAD	C6N-N1N-C2N	-3.43	118.96	121.88
2	B	401	NAD	C6N-N1N-C2N	-2.82	119.48	121.88

There are no chirality outliers.

All (20) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	401	NAD	O4D-C1D-N1N-C6N
2	B	401	NAD	C5D-O5D-PN-O2N
2	B	401	NAD	O4D-C1D-N1N-C6N
2	C	401	NAD	C5B-O5B-PA-O1A
2	C	401	NAD	C5B-O5B-PA-O2A
2	C	401	NAD	C5B-O5B-PA-O3
2	C	401	NAD	O4D-C1D-N1N-C2N
2	C	401	NAD	O4D-C1D-N1N-C6N
2	C	401	NAD	C2D-C1D-N1N-C2N
2	C	401	NAD	C2D-C1D-N1N-C6N
2	C	401	NAD	O4B-C4B-C5B-O5B
2	C	401	NAD	C3B-C4B-C5B-O5B
2	B	401	NAD	O4D-C4D-C5D-O5D
2	B	401	NAD	C3D-C4D-C5D-O5D
2	B	401	NAD	C5D-O5D-PN-O3
2	B	401	NAD	C5D-O5D-PN-O1N
2	A	401	NAD	O4D-C1D-N1N-C2N
2	B	401	NAD	O4D-C1D-N1N-C2N
2	C	401	NAD	PA-O3-PN-O1N
2	C	401	NAD	PA-O3-PN-O2N


There are no ring outliers.


2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	NAD	3	0
2	B	401	NAD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	352/354 (99%)	0.05	4 (1%) 78 74	46, 61, 86, 117	0
1	B	352/354 (99%)	0.08	5 (1%) 73 69	46, 62, 85, 107	0
1	C	352/354 (99%)	1.08	51 (14%) 6 4	55, 89, 126, 151	0
All	All	1056/1062 (99%)	0.40	60 (5%) 29 24	46, 67, 110, 151	0

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	59	LEU	5.5
1	C	30	ASP	5.5
1	A	3	SER	4.2
1	C	246	LEU	4.0
1	C	77	ASN	3.5
1	C	224	ARG	3.3
1	B	94	CYS	3.2
1	C	73	GLN	3.2
1	C	121	GLY	3.1
1	C	175	VAL	3.1
1	C	32	VAL	3.0
1	B	3	SER	3.0
1	C	228	GLY	2.9
1	C	229	ARG	2.9
1	C	240	LYS	2.9
1	C	227	SER	2.9
1	C	354	THR	2.9
1	C	219	LEU	2.8
1	C	218	ALA	2.7
1	A	354	THR	2.7
1	C	176	ALA	2.7
1	B	226	ILE	2.6
1	A	108	CYS	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	3	SER	2.6
1	C	15	ALA	2.5
1	C	92	MET	2.5
1	C	244	LEU	2.5
1	C	49	TYR	2.5
1	C	204	VAL	2.5
1	C	74	ALA	2.5
1	C	88	VAL	2.5
1	C	127	VAL	2.5
1	C	241	ALA	2.5
1	A	229	ARG	2.4
1	C	230	ASP	2.4
1	C	29	ALA	2.4
1	C	81	TRP	2.3
1	C	19	ARG	2.3
1	C	105	GLY	2.3
1	B	354	THR	2.3
1	C	50	TYR	2.3
1	C	24	ILE	2.2
1	C	131	ARG	2.2
1	C	119	GLN	2.2
1	C	221	LEU	2.2
1	C	118	ILE	2.2
1	C	209	LEU	2.2
1	C	226	ILE	2.2
1	C	128	VAL	2.1
1	C	129	PRO	2.1
1	C	31	GLN	2.1
1	C	231	LEU	2.1
1	C	120	GLY	2.1
1	B	84	ARG	2.1
1	C	54	ARG	2.1
1	C	96	THR	2.1
1	C	308	TYR	2.1
1	C	94	CYS	2.0
1	C	143	ALA	2.0
1	C	206	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	B	403	5/5	0.72	0.11	104,110,113,125	0
3	SO4	A	403	5/5	0.88	0.06	97,100,115,115	0
4	ZN	A	405	1/1	0.92	0.09	70,70,70,70	0
4	ZN	C	403	1/1	0.94	0.07	120,120,120,120	0
4	ZN	B	405	1/1	0.95	0.06	73,73,73,73	0
2	NAD	C	401	44/44	0.95	0.09	60,69,112,121	0
2	NAD	A	401	44/44	0.96	0.07	50,61,71,84	0
2	NAD	B	401	44/44	0.96	0.07	50,63,71,78	0
3	SO4	B	402	5/5	0.98	0.07	64,64,71,71	0
3	SO4	A	402	5/5	0.98	0.05	40,46,47,52	0
4	ZN	A	404	1/1	0.98	0.04	56,56,56,56	0
4	ZN	C	402	1/1	1.00	0.03	65,65,65,65	0
4	ZN	B	404	1/1	1.00	0.02	57,57,57,57	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around NAD C 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAD A 401:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.