



# Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2026 – 01:30 pm BST

PDB ID : 9RAK / pdb\_00009rak  
Title : E-selectin complexed with glycomimetic ligand GMI-1077  
Authors : Jakob, R.P.; Ernst, B.  
Deposited on : 2025-05-21  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	280	
1	B	280	

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5292 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called E-selectin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	280	Total	C	N	O	S	0	7	0
			2227	1374	369	456	28			
1	B	280	Total	C	N	O	S	0	5	0
			2216	1369	368	451	28			

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula:  $C_8H_{15}NO_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is (2 {S})-3-cyclohexyl-2-[(2 {R},3 {S},4 {S},5 {R},6 {R})-2-(hydroxymethyl)-6-[(1 {R},2 {R},3 {S})-3-methyl-2-[(2 {S},3 {S},4 {R},5 {S},6 {S})-6-methyl-3,4,5-tris(oxidanyl)oxan-2-yl]oxy-cyclohexyl]oxy-3-oxidanyl-5-(phenylcarbonyloxy)oxan-4-yl]oxy-propanoic acid (CCD ID: A1JDT) (formula: C<sub>35</sub>H<sub>52</sub>O<sub>14</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			49	35	14		
3	B	1	Total	C	O	0	0
			49	35	14		

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total 1	Ca 1	0	0
4	B	1	Total 1	Ca 1	0	0

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total 1	Cl 1	0	0
5	B	1	Total 1	Cl 1	0	0

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	303	Total 303	O 303	0	0
6	B	286	Total 290	O 290	0	4

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: E-selectin

Chain A: 



- Molecule 1: E-selectin

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	51.00Å 58.20Å 65.73Å 96.63° 99.19° 90.45°	Depositor
Resolution (Å)	29.92 – 1.90 29.92 – 1.90	Depositor EDS
% Data completeness (in resolution range)	96.4 (29.92-1.90) 96.4 (29.92-1.90)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.67 (at 1.91Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, $R_{free}$	0.189 , 0.233 0.207 , 0.210	Depositor DCC
$R_{free}$ test set	2701 reflections (4.63%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.2	Xtriage
Anisotropy	0.469	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 50.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5292	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.41% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, A1JDT, CL, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.87	0/2282	1.11	3/3103 (0.1%)
1	B	0.86	0/2271	1.13	2/3089 (0.1%)
All	All	0.86	0/4553	1.12	5/6192 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	124	ASN	CA-CB-CG	7.16	119.76	112.60
1	A	235	PHE	N-CA-C	6.93	119.86	111.82
1	B	273	ASN	CA-CB-CG	6.24	118.84	112.60
1	A	253	GLU	CB-CG-CD	6.20	123.14	112.60
1	B	139	ASN	CA-CB-CG	5.50	118.10	112.60

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2227	0	2029	7	0
1	B	2216	0	2023	22	0
2	A	70	0	65	0	0
2	B	84	0	78	1	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	49	0	0	0	0
3	B	49	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	303	0	0	0	0
6	B	290	0	0	3	0
All	All	5292	0	4195	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:166:GLU:HG2	1:B:167:HIS:CD2	2.36	0.61
1:B:29:ILE:HG12	1:B:104:TRP:HE1	1.66	0.60
1:B:263:LEU:HD22	1:B:273:ASN:HB2	1.87	0.57
1:B:158:ASN:HA	1:B:180:SER:HA	1.89	0.54
1:B:29:ILE:HD11	1:B:93:ILE:HD12	1.90	0.53
1:B:101:VAL:HG23	6:B:641:HOH:O	2.09	0.51
1:A:28:ALA:O	1:A:29:ILE:C	2.52	0.51
1:B:32:LYS:O	1:B:34:GLU:N	2.42	0.50
1:B:157:VAL:HG12	1:B:181:TYR:HB2	1.95	0.49
1:A:29:ILE:HG23	1:A:35:ILE:HD11	1.98	0.46
1:A:158:ASN:HA	1:A:180:SER:HA	1.97	0.46
1:A:157:VAL:HG12	1:A:181:TYR:HB2	1.98	0.46
1:B:33:GLU:HA	1:B:36:GLU:HB2	1.98	0.45
1:B:32:LYS:C	1:B:34:GLU:H	2.23	0.45
1:A:30:GLN:O	1:A:31:ASN:HB2	2.16	0.45
1:B:260:ALA:O	1:B:276:PRO:HG3	2.16	0.44
1:B:27:VAL:HG12	1:B:118:TYR:HB3	1.99	0.44
1:A:62:TRP:CE2	1:A:69:LEU:HG	2.54	0.43
1:B:29:ILE:O	1:B:30:GLN:C	2.60	0.43
1:B:47:SER:OG	6:B:401:HOH:O	2.21	0.43
1:B:29:ILE:HG12	1:B:104:TRP:NE1	2.31	0.42
1:B:230:GLY:HA2	1:B:251:CYS:HA	2.00	0.42
1:B:32:LYS:C	1:B:34:GLU:N	2.79	0.41
1:B:39:ASN:O	1:B:96:LYS:HE3	2.21	0.41
1:B:66:GLN:NE2	6:B:415:HOH:O	2.54	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:31:ASN:CG	1:B:32:LYS:H	2.29	0.40
1:B:161:ALA:HB2	2:B:304:NAG:H62	2.03	0.40
1:A:28:ALA:O	1:A:30:GLN:HG2	2.20	0.40
1:B:249:PHE:HE1	1:B:263:LEU:HG	1.86	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	285/280 (102%)	272 (95%)	11 (4%)	2 (1%)	18	10
1	B	283/280 (101%)	270 (95%)	11 (4%)	2 (1%)	18	10
All	All	568/560 (101%)	542 (95%)	22 (4%)	4 (1%)	18	10

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	29	ILE
1	A	139	ASN
1	B	86	LYS
1	B	139	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	254/247 (103%)	254 (100%)	0	100	100
1	B	252/247 (102%)	250 (99%)	2 (1%)	73	75
All	All	506/494 (102%)	504 (100%)	2 (0%)	84	87

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	29	ILE
1	B	86	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	4	ASN
1	A	31	ASN
1	A	58	ASN
1	A	130	HIS
1	A	237	ASN
1	A	264	GLN
1	A	270	ASN
1	B	58	ASN
1	B	66	GLN
1	B	236	GLN
1	B	237	ASN
1	B	264	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry

Of 17 ligands modelled in this entry, 4 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	A	304	1	14,14,15	0.32	0	17,19,21	1.50	2 (11%)
3	A1JDT	A	306	4	53,53,53	1.30	7 (13%)	69,75,75	1.04	4 (5%)
2	NAG	B	303	1	14,14,15	0.31	0	17,19,21	0.63	1 (5%)
2	NAG	A	305	1	14,14,15	0.30	0	17,19,21	0.46	0
2	NAG	A	301	1	14,14,15	0.32	0	17,19,21	0.81	1 (5%)
2	NAG	B	306	1	14,14,15	0.29	0	17,19,21	0.67	0
2	NAG	B	305	1	14,14,15	0.30	0	17,19,21	0.63	0
2	NAG	B	304	1	14,14,15	0.29	0	17,19,21	0.53	0
2	NAG	B	301	1	14,14,15	0.28	0	17,19,21	0.97	1 (5%)
2	NAG	B	302	1	14,14,15	0.31	0	17,19,21	0.70	1 (5%)
3	A1JDT	B	307	4	53,53,53	1.51	9 (16%)	69,75,75	0.88	3 (4%)
2	NAG	A	302	1	14,14,15	0.27	0	17,19,21	0.70	1 (5%)
2	NAG	A	303	1	14,14,15	0.28	0	17,19,21	0.85	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	304	1	-	1/6/23/26	0/1/1/1
3	A1JDT	A	306	4	-	2/30/92/92	0/5/5/5
2	NAG	B	303	1	-	2/6/23/26	0/1/1/1
2	NAG	A	305	1	-	0/6/23/26	0/1/1/1
2	NAG	A	301	1	-	0/6/23/26	0/1/1/1
2	NAG	B	306	1	-	0/6/23/26	0/1/1/1
2	NAG	B	305	1	-	2/6/23/26	0/1/1/1
2	NAG	B	304	1	-	0/6/23/26	0/1/1/1
2	NAG	B	301	1	-	0/6/23/26	0/1/1/1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	302	1	-	0/6/23/26	0/1/1/1
3	A1JDT	B	307	4	-	1/30/92/92	0/5/5/5
2	NAG	A	302	1	-	0/6/23/26	0/1/1/1
2	NAG	A	303	1	-	2/6/23/26	0/1/1/1

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	307	A1JDT	CAH-CAI	3.73	1.55	1.52
3	B	307	A1JDT	CBS-CBP	-3.57	1.41	1.50
3	A	306	A1JDT	CBS-CBP	-3.53	1.41	1.50
3	B	307	A1JDT	CBJ-CBH	3.00	1.59	1.52
3	A	306	A1JDT	O5-C1	2.75	1.48	1.41
3	B	307	A1JDT	CAE-CAF	2.67	1.60	1.52
3	B	307	A1JDT	CAZ-CBA	2.44	1.59	1.53
3	B	307	A1JDT	CBN-CBL	2.41	1.58	1.52
3	A	306	A1JDT	OBE-CBF	2.32	1.48	1.41
3	A	306	A1JDT	OBO-CBN	-2.29	1.37	1.43
3	A	306	A1JDT	CBB-CBD	2.21	1.58	1.53
3	B	307	A1JDT	O2-C2	-2.09	1.41	1.44
3	A	306	A1JDT	CAY-CAX	2.04	1.56	1.52
3	B	307	A1JDT	O1-CAX	-2.01	1.41	1.44
3	A	306	A1JDT	CBD-CAX	2.01	1.58	1.52
3	B	307	A1JDT	CBD-CAX	2.00	1.58	1.52

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	304	NAG	C1-C2-N2	5.45	119.80	110.49
3	A	306	A1JDT	C1-O1-CAX	-4.00	109.15	115.33
2	B	301	NAG	C1-O5-C5	3.23	116.58	112.19
2	A	301	NAG	C1-O5-C5	2.89	116.10	112.19
2	A	303	NAG	C1-O5-C5	2.73	115.90	112.19
3	A	306	A1JDT	CAG-CAF-CAE	-2.60	106.08	111.73
2	A	302	NAG	C1-O5-C5	2.25	115.24	112.19
3	B	307	A1JDT	OAJ-CAI-CAH	-2.20	117.37	122.57
3	B	307	A1JDT	O1-C1-O5	-2.16	104.63	110.67
3	B	307	A1JDT	C2-O2-CBP	-2.13	113.81	117.21
2	B	302	NAG	C1-O5-C5	2.12	115.06	112.19
3	A	306	A1JDT	C2-O2-CBP	-2.06	113.91	117.21
2	B	303	NAG	C1-O5-C5	2.05	114.97	112.19
2	A	304	NAG	C2-N2-C7	2.01	125.77	122.90

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	306	A1JDT	OAJ-CAI-CAH	-2.01	117.82	122.57

There are no chirality outliers.

All (10) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	303	NAG	C4-C5-C6-O6
2	A	303	NAG	O5-C5-C6-O6
2	B	305	NAG	O5-C5-C6-O6
2	B	303	NAG	O5-C5-C6-O6
3	A	306	A1JDT	O5-C5-C6-O6
3	B	307	A1JDT	O5-C5-C6-O6
2	A	304	NAG	C3-C2-N2-C7
3	A	306	A1JDT	C4-C5-C6-O6
2	A	303	NAG	C4-C5-C6-O6
2	B	305	NAG	C4-C5-C6-O6


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	304	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	280/280 (100%)	0.31	14 (5%)	34 36	10, 27, 47, 80	7 (2%)
1	B	280/280 (100%)	0.41	18 (6%)	25 27	9, 28, 49, 79	5 (1%)
All	All	560/560 (100%)	0.36	32 (5%)	29 31	9, 27, 49, 80	12 (2%)

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	30	GLN	6.0
1	B	235	PHE	5.7
1	B	280	ALA	5.3
1	B	236	GLN	5.0
1	B	137	ILE	4.6
1	A	29	ILE	4.5
1	B	29	ILE	4.5
1	B	237	ASN	4.3
1	B	30	GLN	4.1
1	B	272	ASP	4.0
1	A	32	LYS	4.0
1	A	137	ILE	3.9
1	B	31	ASN	3.9
1	A	31	ASN	3.7
1	A	170	LEU	3.6
1	B	28	ALA	3.5
1	A	253	GLU	3.1
1	B	32	LYS	2.9
1	A	35	ILE	2.8
1	A	238	PRO	2.7
1	B	124[A]	ASN	2.6
1	B	33	GLU	2.5
1	A	237	ASN	2.4
1	B	151	LEU	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	240	SER	2.4
1	B	41	ILE	2.3
1	A	188	SER	2.2
1	A	33	GLU	2.2
1	A	72	GLU	2.1
1	B	72	GLU	2.1
1	B	155	GLN	2.1
1	A	69	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	B	305	14/15	0.52	0.21	79,82,84,84	0
2	NAG	A	302	14/15	0.75	0.14	61,66,69,69	0
2	NAG	B	302	14/15	0.77	0.14	67,69,71,72	0
2	NAG	A	304	14/15	0.77	0.20	94,95,96,96	0
2	NAG	B	306	14/15	0.77	0.13	45,49,56,60	0
2	NAG	A	303	14/15	0.81	0.12	38,43,47,52	0
2	NAG	B	303	14/15	0.82	0.13	39,48,53,57	0
2	NAG	B	304	14/15	0.85	0.13	39,45,49,50	0
2	NAG	B	301	14/15	0.86	0.15	31,42,48,50	0
2	NAG	A	301	14/15	0.90	0.12	31,40,48,52	0
2	NAG	A	305	14/15	0.93	0.07	20,25,28,31	0
3	A1JDT	A	306	49/49	0.95	0.07	14,24,28,32	0
3	A1JDT	B	307	49/49	0.95	0.07	14,23,30,33	0
4	CA	A	307	1/1	0.99	0.02	18,18,18,18	0
4	CA	B	308	1/1	0.99	0.02	17,17,17,17	0

*Continued on next page...*




*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	CL	A	308	1/1	0.99	0.04	21,21,21,21	0
5	CL	B	309	1/1	0.99	0.03	18,18,18,18	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



**Electron density around A1JDT A 306:**

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around A1JDT B 307:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.