Full wwPDB X-ray Structure Validation Report ⓘ

Oct 26, 2026 – 12:53 pm GMT

PDB ID	:	9SZ6 / pdb_00009sz6
Title	:	Crystal structure of the human METTL3-METTL14 in complex with COD_ME_05
Authors	:	Bedi, R.K.; Caflisch, A.
Deposited on	:	2025-10-14
Resolution	:	1.57 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references ⓘ) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.2
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.57 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1094 (1.58-1.58)
Clashscore	190562	1105 (1.58-1.58)
Ramachandran outliers	187476	1082 (1.58-1.58)
Sidechain outliers	187428	1081 (1.58-1.58)
RSRZ outliers	180081	1094 (1.58-1.58)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	228	11% 85% 5% 10%
2	B	307	11% 73% 7% 20%

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 3864 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N6-adenosine-methyltransferase catalytic subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	205	1609	1033	280	287	9	0	1	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	353	MET	-	initiating methionine	UNP Q86U44

- Molecule 2 is a protein called N(6)-adenosine-methyltransferase non-catalytic subunit METTL14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	247	1900	1218	324	345	13	0	1	0

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	89	MET	-	initiating methionine	UNP Q9HCE5
B	90	HIS	-	expression tag	UNP Q9HCE5
B	91	HIS	-	expression tag	UNP Q9HCE5
B	92	HIS	-	expression tag	UNP Q9HCE5
B	93	HIS	-	expression tag	UNP Q9HCE5
B	94	HIS	-	expression tag	UNP Q9HCE5
B	95	HIS	-	expression tag	UNP Q9HCE5
B	96	SER	-	expression tag	UNP Q9HCE5
B	97	SER	-	expression tag	UNP Q9HCE5
B	98	GLY	-	expression tag	UNP Q9HCE5
B	99	ARG	-	expression tag	UNP Q9HCE5
B	100	GLU	-	expression tag	UNP Q9HCE5
B	101	ASN	-	expression tag	UNP Q9HCE5
B	102	LEU	-	expression tag	UNP Q9HCE5

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	103	TYR	-	expression tag	UNP Q9HCE5
B	104	PHE	-	expression tag	UNP Q9HCE5
B	105	GLN	-	expression tag	UNP Q9HCE5
B	106	GLY	-	expression tag	UNP Q9HCE5

- Molecule 3 is 2-[[6-[[(3-fluoranyl-1-bicyclo[1.1.1]pentanyl)methylamino]methyl]imidazo[1,2-a]pyridin-2-yl]methyl]-5-(7-oxa-2-azaspiro[3.5]nonan-2-yl)-2,7-naphthyridin-1-one (CCD ID: A1JSB) (formula: $C_{30}H_{33}FN_6O_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	F	N	O	0	0
			39	30	1	6	2		

- Molecule 4 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C O 4 2 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	134	Total O 134 134	0	0
5	B	178	Total O 178 178	0	0


3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: N6-adenosine-methyltransferase catalytic subunit



• Molecule 2: N(6)-adenosine-methyltransferase non-catalytic subunit METTL14



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α, β, γ	63.98Å 63.98Å 225.74Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	35.00 – 1.57 35.00 – 1.57	Depositor EDS
% Data completeness (in resolution range)	99.9 (35.00-1.57) 99.8 (35.00-1.57)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.15 (at 1.57Å)	Xtriage
Refinement program	PHENIX 1.21.1_5286	Depositor
R, R_{free}	0.195 , 0.203 0.195 , 0.203	Depositor DCC
R_{free} test set	1994 reflections (2.62%)	wwPDB-VP
Wilson B-factor (Å^2)	30.2	Xtriage
Anisotropy	0.120	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.36 , 44.7	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.025 for -h,-k,l	Xtriage
F_o,F_c correlation	0.97	EDS
Total number of atoms	3864	wwPDB-VP
Average B, all atoms (Å^2)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.21% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: A1JSB, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	A	0.26	0/1651	0.52	0/2252
2	B	0.28	0/1951	0.50	0/2655
All	All	0.27	0/3602	0.51	0/4907

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1609	0	1525	7	0
2	B	1900	0	1754	13	0
3	A	39	0	0	0	0
4	B	4	0	3	0	0
5	A	134	0	0	2	0
5	B	178	0	0	2	0
All	All	3864	0	3282	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:490:ASN:ND2	5:A:701:HOH:O	2.05	0.90
2:B:254:ARG:NH2	2:B:294:GLY:O	2.17	0.78
2:B:168:MET:HE3	2:B:369:TYR:HA	1.75	0.68
2:B:180:ARG:NH2	2:B:220:GLU:OE1	2.33	0.61
2:B:152:LEU:O	2:B:154:ARG:N	2.37	0.58
2:B:198:TYR:HB3	2:B:324:ILE:HG22	1.91	0.53
1:A:474:HIS:O	2:B:307:ALA:N	2.44	0.51
1:A:433[A]:THR:HG23	5:A:814:HOH:O	2.10	0.51
2:B:222:ASP:OD2	5:B:501:HOH:O	2.19	0.47
2:B:178:ASP:OD1	2:B:180:ARG:NH1	2.48	0.47
1:A:539:ASN:O	1:A:541:GLN:HG2	2.15	0.47
1:A:442:GLU:O	1:A:446:LEU:HG	2.16	0.46
2:B:220:GLU:OE2	5:B:502:HOH:O	2.21	0.45
2:B:179:ILE:HG12	2:B:224:ILE:HD13	1.99	0.45
1:A:373:TRP:HB2	1:A:551:LEU:HD13	2.00	0.43
2:B:311:ILE:HG23	2:B:313:LEU:H	1.84	0.42
1:A:391:VAL:HB	1:A:530:LYS:HG2	2.01	0.42
2:B:168:MET:HE1	2:B:374:TYR:HB2	2.02	0.41
2:B:260[B]:CYS:SG	2:B:285:LYS:HD2	2.61	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	202/228 (89%)	197 (98%)	4 (2%)	1 (0%)	24 9
2	B	240/307 (78%)	233 (97%)	6 (2%)	1 (0%)	30 12
All	All	442/535 (83%)	430 (97%)	10 (2%)	2 (0%)	24 9

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	467	ILE
2	B	153	ILE

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	165/200 (82%)	165 (100%)	0	100	100
2	B	189/274 (69%)	187 (99%)	2 (1%)	65	43
All	All	354/474 (75%)	352 (99%)	2 (1%)	78	65

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	217	MET
2	B	311	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	464	GLN
1	A	567	GLN
2	B	122	HIS
2	B	282	GLN
2	B	382	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	A1JSB	A	601	-	44,46,46	2.89	24 (54%)	38,72,72	2.36	8 (21%)
4	ACT	B	401	-	3,3,3	1.19	0	3,3,3	0.97	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	A1JSB	A	601	-	-	2/13/53/53	0/9/8/8

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	601	A1JSB	C17-C07	6.53	1.48	1.41
3	A	601	A1JSB	C35-C32	6.03	1.60	1.53
3	A	601	A1JSB	C34-C32	5.87	1.60	1.53
3	A	601	A1JSB	C31-C32	5.42	1.59	1.53
3	A	601	A1JSB	C23-C22	4.30	1.43	1.36
3	A	601	A1JSB	C29-N28	3.71	1.51	1.46
3	A	601	A1JSB	C29-C30	-3.69	1.49	1.53
3	A	601	A1JSB	C37-C38	3.66	1.48	1.41
3	A	601	A1JSB	C16-N08	3.63	1.53	1.48
3	A	601	A1JSB	C31-C30	3.61	1.59	1.55
3	A	601	A1JSB	C38-N24	3.60	1.42	1.39
3	A	601	A1JSB	C09-N08	3.57	1.53	1.48

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	601	A1JSB	F33-C32	-3.42	1.32	1.42
3	A	601	A1JSB	C23-N24	3.32	1.42	1.37
3	A	601	A1JSB	C19-N20	3.20	1.41	1.37
3	A	601	A1JSB	C19-C18	3.00	1.40	1.33
3	A	601	A1JSB	C04-C03	2.84	1.45	1.40
3	A	601	A1JSB	C06-N05	2.54	1.39	1.34
3	A	601	A1JSB	C21-C22	2.53	1.53	1.50
3	A	601	A1JSB	C34-C30	2.53	1.58	1.55
3	A	601	A1JSB	C06-C07	2.48	1.42	1.39
3	A	601	A1JSB	C38-N39	2.47	1.37	1.33
3	A	601	A1JSB	C17-C18	2.06	1.49	1.44
3	A	601	A1JSB	C15-C14	2.03	1.57	1.51

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	A	601	A1JSB	C03-C02-N20	10.71	120.05	113.80
3	A	601	A1JSB	C23-N24-C25	3.52	133.42	129.98
3	A	601	A1JSB	C19-N20-C02	-2.96	120.69	122.88
3	A	601	A1JSB	N24-C38-N39	-2.91	108.96	110.61
3	A	601	A1JSB	C37-C36-C26	-2.85	118.41	121.61
3	A	601	A1JSB	C04-C03-C17	2.35	120.60	117.84
3	A	601	A1JSB	C14-O13-C12	2.32	117.64	109.89
3	A	601	A1JSB	C17-C03-C02	-2.31	120.00	121.63

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	601	A1JSB	C22-C21-N20-C02
3	A	601	A1JSB	C22-C21-N20-C19


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring

in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand A1JSB A 601



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	205/228 (89%)	0.62	24 (11%)	9	15	15, 33, 58, 65	1 (0%)
2	B	247/307 (80%)	0.78	33 (13%)	7	12	17, 32, 63, 80	1 (0%)
All	All	452/535 (84%)	0.71	57 (12%)	8	13	15, 33, 61, 80	2 (0%)

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	307	ALA	6.6
2	B	137	VAL	5.3
2	B	155	LEU	5.0
2	B	308	ASN	5.0
1	A	511	SER	4.8
2	B	153	ILE	4.8
2	B	202	THR	4.5
1	A	573	ILE	4.4
2	B	392	ILE	4.3
1	A	368	LEU	4.3
2	B	274	THR	4.3
2	B	309	VAL	4.2
1	A	467	ILE	4.1
2	B	152	LEU	3.9
2	B	295	THR	3.8
2	B	270	GLY	3.7
2	B	272	THR	3.7
2	B	321	ILE	3.6
1	A	575	SER	3.6
2	B	275	LEU	3.3
1	A	509	SER	3.3
1	A	576	LYS	3.2
2	B	117	ASN	3.2
1	A	468	ARG	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	177	PHE	3.1
1	A	446	LEU	3.0
1	A	463	LEU	3.0
1	A	475	TRP	3.0
1	A	512	HIS	2.9
2	B	179	ILE	2.9
1	A	367	ARG	2.8
2	B	273	LYS	2.8
1	A	402	MET	2.8
1	A	508	ARG	2.8
1	A	574	ILE	2.7
1	A	510	THR	2.7
1	A	474	HIS	2.7
2	B	393	GLU	2.7
1	A	507	VAL	2.6
2	B	388	CYS	2.6
2	B	176	ALA	2.6
2	B	151	GLU	2.5
1	A	490	ASN	2.5
1	A	462	GLN	2.4
2	B	390	GLU	2.4
2	B	387	GLY	2.3
2	B	210	CYS	2.3
2	B	389	THR	2.3
1	A	477	ASN	2.2
2	B	271	LYS	2.2
1	A	405	PRO	2.2
2	B	391	GLU	2.1
2	B	380	ALA	2.1
2	B	154	ARG	2.1
1	A	369	PHE	2.0
2	B	322	GLY	2.0
2	B	381	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	A1JSB	A	601	39/39	0.94	0.09	27,31,42,46	0
4	ACT	B	401	4/4	0.96	0.07	28,28,29,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around A1JSB A 601:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers

There are no such residues in this entry.