



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 19, 2026 – 01:38 pm BST

PDB ID : 9SZF / pdb_00009szf
Title : Crystal structure of the human METTL3-METTL14 in complex with
COD_ME_27
Authors : Bedi, R.K.; Caflisch, A.
Deposited on : 2025-10-14
Resolution : 1.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.2
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1112 (1.82-1.82)
Clashscore	190562	1148 (1.82-1.82)
Ramachandran outliers	187476	1140 (1.82-1.82)
Sidechain outliers	187428	1140 (1.82-1.82)
RSRZ outliers	180081	1112 (1.82-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	228	11% 78% 10% 13%
2	B	307	10% 72% 6% 22%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3641 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N6-adenosine-methyltransferase catalytic subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	199	Total	C	N	O	S	0	0	0
			1553	997	272	275	9			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	353	MET	-	initiating methionine	UNP Q86U44

- Molecule 2 is a protein called N(6)-adenosine-methyltransferase non-catalytic subunit METTL14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	240	Total	C	N	O	S	0	1	0
			1845	1185	314	333	13			

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	89	MET	-	initiating methionine	UNP Q9HCE5
B	90	HIS	-	expression tag	UNP Q9HCE5
B	91	HIS	-	expression tag	UNP Q9HCE5
B	92	HIS	-	expression tag	UNP Q9HCE5
B	93	HIS	-	expression tag	UNP Q9HCE5
B	94	HIS	-	expression tag	UNP Q9HCE5
B	95	HIS	-	expression tag	UNP Q9HCE5
B	96	SER	-	expression tag	UNP Q9HCE5
B	97	SER	-	expression tag	UNP Q9HCE5
B	98	GLY	-	expression tag	UNP Q9HCE5
B	99	ARG	-	expression tag	UNP Q9HCE5
B	100	GLU	-	expression tag	UNP Q9HCE5
B	101	ASN	-	expression tag	UNP Q9HCE5
B	102	LEU	-	expression tag	UNP Q9HCE5


Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	103	TYR	-	expression tag	UNP Q9HCE5
B	104	PHE	-	expression tag	UNP Q9HCE5
B	105	GLN	-	expression tag	UNP Q9HCE5
B	106	GLY	-	expression tag	UNP Q9HCE5

- Molecule 3 is {N}-(2-methylpropyl)-2-[4-[[5-(7-oxa-2-azaspiro[3.5]nonan-2-yl)-1-oxidanylidene-2,7-naphthyridin-2-yl]methyl]phenyl]- {N}-[[2 {S}]-piperidin-2-yl]methyl]ethanamide (CCD ID: A1JSD) (formula: C₃₄H₄₅N₅O₃) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	89	Total	O	0	0
			89	89		
5	B	108	Total	O	0	0
			108	108		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: N6-adenosine-methyltransferase catalytic subunit



- Molecule 2: N(6)-adenosine-methyltransferase non-catalytic subunit METTL14



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	64.14Å 64.14Å 225.12Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	44.65 – 1.82 44.65 – 1.82	Depositor EDS
% Data completeness (in resolution range)	99.9 (44.65-1.82) 99.9 (44.65-1.82)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.30 (at 1.82Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.197 , 0.216 0.198 , 0.214	Depositor DCC
R_{free} test set	2423 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	37.3	Xtriage
Anisotropy	0.077	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 44.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.021 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3641	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, A1JSD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/1592	0.57	0/2168
2	B	0.35	0/1896	0.56	0/2581
All	All	0.34	0/3488	0.56	0/4749

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	B	0	1
All	All	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	536	ARG	Sidechain
2	B	180	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1553	0	1477	13	0
2	B	1845	0	1699	10	0
3	A	42	0	0	0	0
4	B	4	0	3	0	0
5	A	89	0	0	3	0
5	B	108	0	0	1	0
All	All	3641	0	3179	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:222:ASP:OD1	5:B:501:HOH:O	1.90	0.88
2:B:168:MET:HE3	2:B:369:TYR:HA	1.74	0.70
1:A:373:TRP:HB2	1:A:551:LEU:HD13	1.78	0.65
1:A:539:ASN:O	1:A:541:GLN:HG2	2.03	0.58
1:A:391:VAL:HB	1:A:530:LYS:HG2	1.84	0.57
1:A:488:LYS:HE3	5:A:715:HOH:O	2.03	0.57
2:B:311:ILE:HG23	2:B:313:LEU:H	1.71	0.55
2:B:263:LYS:HD3	2:B:318:GLU:HG3	1.89	0.54
2:B:198:TYR:HB3	2:B:324:ILE:HG22	1.92	0.52
1:A:533:LEU:CD2	1:A:547:LEU:HD12	2.44	0.47
2:B:184:PRO:HG2	2:B:342:ARG:HD3	1.96	0.46
2:B:214:ASP:O	2:B:218:LYS:HG3	2.16	0.44
1:A:539:ASN:ND2	5:A:706:HOH:O	2.46	0.44
1:A:423:GLN:O	1:A:488:LYS:NZ	2.47	0.43
1:A:456:ILE:HD13	2:B:260[B]:CYS:SG	2.59	0.42
2:B:260[B]:CYS:SG	2:B:285:LYS:HD2	2.60	0.41
1:A:555:HIS:HE1	5:A:723:HOH:O	2.04	0.41
1:A:533:LEU:HD22	1:A:547:LEU:HD12	2.02	0.41
1:A:412:ASP:O	1:A:416:ARG:HG3	2.21	0.41
2:B:168:MET:HE1	2:B:374:TYR:HB2	2.03	0.41
1:A:418:ASN:O	1:A:421:VAL:HG12	2.21	0.40
1:A:457:TRP:CE3	1:A:514:PRO:HD3	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	193/228 (85%)	188 (97%)	4 (2%)	1 (0%)	24	14
2	B	233/307 (76%)	227 (97%)	5 (2%)	1 (0%)	30	19
All	All	426/535 (80%)	415 (97%)	9 (2%)	2 (0%)	24	14

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	467	ILE
2	B	152	LEU

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	159/200 (80%)	157 (99%)	2 (1%)	61	50
2	B	183/274 (67%)	183 (100%)	0	100	100
All	All	342/474 (72%)	340 (99%)	2 (1%)	78	74

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	462	GLN
1	A	507	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	512	HIS
1	A	541	GLN
2	B	122	HIS
2	B	171	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	A1JSD	A	601	-	47,47,47	2.79	12 (25%)	44,67,67	2.72	11 (25%)
4	ACT	B	401	-	3,3,3	1.44	1 (33%)	3,3,3	1.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	A1JSD	A	601	-	-	0/24/52/52	0/6/6/6

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	601	A1JSD	C24-C25	12.14	1.53	1.41
3	A	601	A1JSD	C22-N21	6.16	1.46	1.37
3	A	601	A1JSD	C20-N21	5.14	1.56	1.47
3	A	601	A1JSD	C24-C23	4.54	1.54	1.44
3	A	601	A1JSD	C34-N26	3.68	1.53	1.48
3	A	601	A1JSD	C35-N36	3.54	1.41	1.34
3	A	601	A1JSD	C37-C38	3.48	1.47	1.40
3	A	601	A1JSD	C38-C39	3.44	1.54	1.47
3	A	601	A1JSD	C15-C16	3.29	1.57	1.51
3	A	601	A1JSD	C25-N26	3.03	1.47	1.41
3	A	601	A1JSD	C27-N26	2.97	1.52	1.48
3	A	601	A1JSD	C39-N21	2.59	1.42	1.39
4	B	401	ACT	CH3-C	2.06	1.57	1.49

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	601	A1JSD	C38-C39-N21	13.51	121.68	113.80
3	A	601	A1JSD	C22-N21-C39	-5.34	118.94	122.88
3	A	601	A1JSD	C24-C38-C39	-4.43	118.52	121.63
3	A	601	A1JSD	C19-C20-N21	-4.29	106.35	112.83
3	A	601	A1JSD	C37-N36-C35	3.03	121.61	117.48
3	A	601	A1JSD	C37-C38-C24	2.79	121.11	117.84
3	A	601	A1JSD	O14-C13-C15	-2.72	117.18	121.60
3	A	601	A1JSD	C15-C13-N05	2.55	122.17	117.74
3	A	601	A1JSD	C16-C15-C13	2.54	117.08	112.61
3	A	601	A1JSD	C32-O31-C30	2.26	117.42	109.89
3	A	601	A1JSD	C23-C22-N21	-2.20	120.36	122.43

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	199/228 (87%)	0.58	24 (12%)	8 9	28, 41, 70, 76	0
2	B	240/307 (78%)	0.72	32 (13%)	7 7	21, 40, 76, 91	1 (0%)
All	All	439/535 (82%)	0.66	56 (12%)	7 8	21, 41, 72, 91	1 (0%)

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	137	VAL	5.0
2	B	153	ILE	4.6
1	A	475	TRP	4.5
2	B	272	THR	4.3
1	A	368	LEU	4.2
2	B	117	ASN	4.1
1	A	467	ILE	4.1
2	B	270	GLY	3.8
2	B	154	ARG	3.8
1	A	573	ILE	3.8
2	B	152	LEU	3.7
2	B	390	GLU	3.7
1	A	575	SER	3.7
1	A	576	LYS	3.5
1	A	474	HIS	3.4
2	B	151	GLU	3.4
2	B	274	THR	3.4
2	B	118	ASP	3.4
2	B	310	ASP	3.3
2	B	209	LYS	3.2
1	A	463	LEU	3.1
1	A	511	SER	3.1
2	B	275	LEU	3.1
2	B	119	TYR	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	570	PRO	3.0
1	A	367	ARG	3.0
1	A	507	VAL	2.9
2	B	273	LYS	2.9
1	A	407	GLY	2.9
2	B	124	VAL	2.9
2	B	311	ILE	2.9
2	B	321	ILE	2.7
2	B	309	VAL	2.7
2	B	155	LEU	2.6
1	A	468	ARG	2.6
2	B	199	TYR	2.5
2	B	322	GLY	2.5
2	B	271	LYS	2.5
2	B	120	CYS	2.4
2	B	389	THR	2.4
1	A	574	ILE	2.4
1	A	478	HIS	2.4
2	B	294	GLY	2.3
2	B	176	ALA	2.3
1	A	466	ILE	2.3
2	B	268	ASN	2.3
1	A	477	ASN	2.2
2	B	200	ARG	2.2
1	A	369	PHE	2.1
2	B	269	PRO	2.1
1	A	462	GLN	2.1
1	A	400	ILE	2.1
1	A	509	SER	2.1
1	A	446	LEU	2.1
2	B	384	TYR	2.0
1	A	494	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	A1JSD	A	601	42/42	0.94	0.09	33,38,46,56	0
4	ACT	B	401	4/4	0.97	0.07	32,33,34,34	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.