



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 21, 2026 – 12:36 PM JST

PDB ID : 9UM7 / pdb_00009um7
Title : CaPETaseM9 SEC loop of 12CL variant
Authors : Kim, K.; Ki, D.; Park, J.
Deposited on : 2025-04-21
Resolution : 1.26 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.26 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1693 (1.28-1.24)
Clashscore	190562	1730 (1.28-1.24)
Ramachandran outliers	187476	1695 (1.28-1.24)
Sidechain outliers	187428	1694 (1.28-1.24)
RSRZ outliers	180081	1693 (1.28-1.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	266	<div> <div>8%</div> <div>91%</div> <div>6% ..</div> </div>
1	B	266	<div> <div>8%</div> <div>92%</div> <div>6% .</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4401 atoms, of which 18 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cutinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	264	Total	C	N	O	S	0	0	0
			2019	1276	356	377	10			
1	B	260	Total	C	N	O	S	0	0	0
			1980	1253	346	371	10			

There are 50 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	109	ALA	ASN	engineered mutation	UNP A0A1M7II12
A	129	THR	VAL	engineered mutation	UNP A0A1M7II12
A	155	ARG	ALA	engineered mutation	UNP A0A1M7II12
A	180	CYS	LEU	engineered mutation	UNP A0A1M7II12
A	196	THR	GLY	engineered mutation	UNP A0A1M7II12
A	198	LYS	ARG	engineered mutation	UNP A0A1M7II12
A	202	CYS	ALA	engineered mutation	UNP A0A1M7II12
A	242	CYS	ARG	engineered mutation	UNP A0A1M7II12
A	280	GLY	-	insertion	UNP A0A1M7II12
A	281	ALA	PRO	engineered mutation	UNP A0A1M7II12
A	282	GLU	GLY	engineered mutation	UNP A0A1M7II12
A	283	HIS	PRO	engineered mutation	UNP A0A1M7II12
A	284	GLN	SER	engineered mutation	UNP A0A1M7II12
A	285	ALA	THR	engineered mutation	UNP A0A1M7II12
A	286	ASP	GLY	engineered mutation	UNP A0A1M7II12
A	288	SER	-	insertion	UNP A0A1M7II12
A	289	LYS	PHE	engineered mutation	UNP A0A1M7II12
A	290	ARG	ALA	engineered mutation	UNP A0A1M7II12
A	291	ALA	PRO	engineered mutation	UNP A0A1M7II12
A	293	CYS	SER	engineered mutation	UNP A0A1M7II12
A	302	LEU	-	expression tag	UNP A0A1M7II12
A	303	GLU	-	expression tag	UNP A0A1M7II12
A	304	HIS	-	expression tag	UNP A0A1M7II12
A	305	HIS	-	expression tag	UNP A0A1M7II12
A	306	HIS	-	expression tag	UNP A0A1M7II12

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	109	ALA	ASN	engineered mutation	UNP A0A1M7II12
B	129	THR	VAL	engineered mutation	UNP A0A1M7II12
B	155	ARG	ALA	engineered mutation	UNP A0A1M7II12
B	180	CYS	LEU	engineered mutation	UNP A0A1M7II12
B	196	THR	GLY	engineered mutation	UNP A0A1M7II12
B	198	LYS	ARG	engineered mutation	UNP A0A1M7II12
B	202	CYS	ALA	engineered mutation	UNP A0A1M7II12
B	242	CYS	ARG	engineered mutation	UNP A0A1M7II12
B	280	GLY	-	insertion	UNP A0A1M7II12
B	281	ALA	PRO	engineered mutation	UNP A0A1M7II12
B	282	GLU	GLY	engineered mutation	UNP A0A1M7II12
B	283	HIS	PRO	engineered mutation	UNP A0A1M7II12
B	284	GLN	SER	engineered mutation	UNP A0A1M7II12
B	285	ALA	THR	engineered mutation	UNP A0A1M7II12
B	286	ASP	GLY	engineered mutation	UNP A0A1M7II12
B	288	SER	-	insertion	UNP A0A1M7II12
B	289	LYS	PHE	engineered mutation	UNP A0A1M7II12
B	290	ARG	ALA	engineered mutation	UNP A0A1M7II12
B	291	ALA	PRO	engineered mutation	UNP A0A1M7II12
B	293	CYS	SER	engineered mutation	UNP A0A1M7II12
B	302	LEU	-	expression tag	UNP A0A1M7II12
B	303	GLU	-	expression tag	UNP A0A1M7II12
B	304	HIS	-	expression tag	UNP A0A1M7II12
B	305	HIS	-	expression tag	UNP A0A1M7II12
B	306	HIS	-	expression tag	UNP A0A1M7II12

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C H O 10 2 6 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C H O 10 2 6 2	0	0
2	B	1	Total C H O 10 2 6 2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	176	Total O 176 176	0	0
3	B	184	Total O 184 184	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cutinase



- Molecule 1: Cutinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	77.95Å 77.95Å 350.63Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.68 – 1.26 48.68 – 1.26	Depositor EDS
% Data completeness (in resolution range)	94.9 (48.68-1.26) 95.0 (48.68-1.26)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.04 (at 1.26Å)	Xtriage
Refinement program	REFMAC 5.8.0425	Depositor
R, R_{free}	0.165 , 0.184 0.174 , 0.190	Depositor DCC
R_{free} test set	8084 reflections (4.77%)	wwPDB-VP
Wilson B-factor (Å ²)	17.0	Xtriage
Anisotropy	0.008	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 30.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4401	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.89	3/2078 (0.1%)	1.20	10/2839 (0.4%)
1	B	0.85	0/2036	1.16	6/2782 (0.2%)
All	All	0.87	3/4114 (0.1%)	1.18	16/5621 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	2
All	All	0	5

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	56	SER	CA-CB	5.99	1.62	1.53
1	A	192	ALA	CA-C	-5.80	1.49	1.53
1	A	192	ALA	C-O	5.12	1.26	1.23

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	182	ARG	CD-NE-CZ	6.91	134.08	124.40
1	B	99	PRO	O-C-N	6.60	131.04	122.86
1	B	182	ARG	CD-NE-CZ	-6.50	115.30	124.40
1	B	282	GLU	CB-CG-CD	6.42	123.51	112.60
1	A	306	HIS	CB-CG-CD2	-6.37	122.92	131.20
1	B	237	ASP	CA-CB-CG	6.22	118.82	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	282	GLU	O-C-N	6.12	129.80	122.27
1	A	269	ASP	CA-CB-CG	6.02	118.62	112.60
1	B	232	THR	CA-CB-OG1	-5.62	101.17	109.60
1	A	247	PHE	CA-CB-CG	-5.43	108.37	113.80
1	A	99	PRO	CA-C-N	5.31	131.82	121.41
1	A	99	PRO	C-N-CA	5.31	131.82	121.41
1	A	248	PHE	CA-CB-CG	5.31	119.11	113.80
1	A	186	LYS	CD-CE-NZ	5.27	128.77	111.90
1	A	282	GLU	CB-CG-CD	5.22	121.48	112.60
1	B	254	THR	CA-CB-OG1	-5.13	101.91	109.60

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	182	ARG	Sidechain
1	A	99	PRO	Peptide,Mainchain
1	B	155	ARG	Sidechain
1	B	182	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2019	0	1931	11	0
1	B	1980	0	1904	3	0
2	A	8	6	12	2	0
2	B	16	12	24	0	0
3	A	176	0	0	1	1
3	B	184	0	0	0	0
All	All	4383	18	3871	14	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:235:GLU:H	1:A:306:HIS:HE1	1.23	0.87
1:A:216:ALA:H	2:A:402:EDO:H22	1.40	0.85
1:B:102:ILE:HA	1:B:130:ILE:HG12	1.82	0.61
1:A:100:GLY:HA2	1:A:168:TRP:O	2.00	0.60
1:A:186:LYS:HD3	3:A:664:HOH:O	2.05	0.56
1:A:100:GLY:HA3	1:A:168:TRP:CE3	2.41	0.55
1:B:101:PHE:HB2	1:B:170:MET:HG3	1.93	0.51
1:A:50:PRO:HD2	1:A:63:PRO:HG3	1.97	0.47
1:A:107:GLN:HE21	1:A:107:GLN:HA	1.80	0.46
1:A:165:ALA:O	1:A:188:ILE:HA	2.16	0.44
1:A:186:LYS:HE2	1:A:267:TRP:O	2.17	0.44
1:A:247:PHE:HB2	2:A:401:EDO:H12	1.99	0.43
1:A:235:GLU:H	1:A:306:HIS:CE1	2.15	0.41
1:B:165:ALA:O	1:B:188:ILE:HA	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:628:HOH:O	3:A:628:HOH:O[12_565]	2.14	0.06

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	262/266 (98%)	253 (97%)	9 (3%)	0	100	100
1	B	258/266 (97%)	253 (98%)	4 (2%)	1 (0%)	30	9
All	All	520/532 (98%)	506 (97%)	13 (2%)	1 (0%)	43	16

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	301	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	210/210 (100%)	206 (98%)	4 (2%)	50	13
1	B	206/210 (98%)	204 (99%)	2 (1%)	68	34
All	All	416/420 (99%)	410 (99%)	6 (1%)	59	22

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	102	ILE
1	A	107	GLN
1	A	293	CYS
1	A	305	HIS
1	B	129	THR
1	B	293	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	81	GLN
1	A	107	GLN
1	A	305	HIS
1	A	306	HIS
1	B	54	ASN
1	B	81	GLN
1	B	181	GLN
1	B	276	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	A	401	-	3,3,3	0.10	0	2,2,2	0.18	0
2	EDO	B	401	-	3,3,3	0.22	0	2,2,2	0.37	0
2	EDO	A	402	-	3,3,3	1.03	0	2,2,2	0.72	0
2	EDO	B	404	-	3,3,3	0.17	0	2,2,2	0.05	0
2	EDO	B	402	-	3,3,3	0.49	0	2,2,2	0.53	0
2	EDO	B	403	-	3,3,3	0.04	0	2,2,2	0.16	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	401	-	-	1/1/1/1	-
2	EDO	B	401	-	-	0/1/1/1	-
2	EDO	A	402	-	-	1/1/1/1	-
2	EDO	B	404	-	-	0/1/1/1	-
2	EDO	B	402	-	-	0/1/1/1	-
2	EDO	B	403	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	402	EDO	O1-C1-C2-O2
2	A	401	EDO	O1-C1-C2-O2




There are no ring outliers.


2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	EDO	1	0
2	A	402	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	264/266 (99%)	0.40	21 (7%) 18 20	12, 17, 28, 55	0
1	B	260/266 (97%)	0.29	20 (7%) 19 21	11, 16, 29, 57	0
All	All	524/532 (98%)	0.35	41 (7%) 19 21	11, 17, 29, 57	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	302	LEU	6.4
1	B	291	ALA	6.0
1	B	293	CYS	5.7
1	B	101	PHE	5.4
1	A	101	PHE	5.0
1	A	305	HIS	4.6
1	A	293	CYS	3.9
1	B	202	CYS	3.7
1	B	43	ASP	3.6
1	A	281	ALA	3.5
1	A	306	HIS	3.2
1	A	73	GLY	3.2
1	A	43	ASP	3.2
1	B	288	SER	3.1
1	A	100	GLY	3.0
1	A	74	ALA	3.0
1	A	151	ARG	3.0
1	A	75	SER	3.0
1	B	75	SER	2.8
1	A	102	ILE	2.7
1	A	72	VAL	2.7
1	B	151	ARG	2.6
1	B	243	ASN	2.6
1	B	73	GLY	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	292	VAL	2.5
1	A	82	ILE	2.5
1	B	290	ARG	2.5
1	A	180	CYS	2.5
1	A	247	PHE	2.4
1	B	301	PHE	2.4
1	B	282	GLU	2.3
1	B	242	CYS	2.3
1	A	290	ARG	2.3
1	A	299	CYS	2.2
1	A	280	GLY	2.2
1	B	247	PHE	2.2
1	A	196	THR	2.2
1	B	150	THR	2.2
1	B	72	VAL	2.1
1	B	100	GLY	2.1
1	A	242	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ


There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	EDO	A	401	4/4	0.65	0.17	20,20,20,20	0
2	EDO	B	404	4/4	0.68	0.17	20,20,20,20	0
2	EDO	B	402	4/4	0.72	0.17	26,27,27,39	0
2	EDO	B	403	4/4	0.74	0.16	20,20,20,20	0
2	EDO	A	402	4/4	0.82	0.21	22,25,27,52	0
2	EDO	B	401	4/4	0.91	0.12	24,25,26,32	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around EDO B 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around EDO B 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around EDO B 403:


$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around EDO A 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.