Full wwPDB X-ray Structure Validation Report (i)

Oct 25, 2026 – 10:09 AM EDT

PDB ID	:	9YXS / pdb_00009yxs
Title	:	Crystal structure of M. tuberculosis ProRS bound to compound 5a
Authors	:	Gade, P.; Michalska, K.; Joachimiak, A.; Center for Structural Biology of Infectious Diseases (CSBID)
Deposited on	:	2025-10-27
Resolution	:	2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.2
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	588	7% 89% 6% .
1	B	588	7% 82% 12% 6%

2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 8706 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Proline–tRNA ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	564	Total	C	N	O	S	0	0	0
			4295	2717	759	811	8			
1	B	552	Total	C	N	O	S	0	0	0
			4217	2664	749	796	8			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	583	GLU	-	expression tag	UNP P9WFT9
A	584	ASN	-	expression tag	UNP P9WFT9
A	585	LEU	-	expression tag	UNP P9WFT9
A	586	TYR	-	expression tag	UNP P9WFT9
A	587	PHE	-	expression tag	UNP P9WFT9
A	588	GLN	-	expression tag	UNP P9WFT9
B	583	GLU	-	expression tag	UNP P9WFT9
B	584	ASN	-	expression tag	UNP P9WFT9
B	585	LEU	-	expression tag	UNP P9WFT9
B	586	TYR	-	expression tag	UNP P9WFT9
B	587	PHE	-	expression tag	UNP P9WFT9
B	588	GLN	-	expression tag	UNP P9WFT9

- Molecule 2 is 3-(4-chloro-2-fluorobenzamido)pyrazine-2-carboxamide (CCD ID: A1CZG) (formula: $C_{12}H_8ClFN_4O_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C Cl F N O 20 12 1 1 4 2	0	0
2	B	1	Total C Cl F N O 20 12 1 1 4 2	0	0


- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	2	Total Mg 2 2	0	0
3	B	1	Total Mg 1 1	0	0

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Ca 1 1	0	0

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0

- Molecule 6 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 4 2 2	0	0

- Molecule 7 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	83	Total O 83 83	0	0
7	B	55	Total O 55 55	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Proline–tRNA ligase

Chain A:



• Molecule 1: Proline–tRNA ligase

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	86.01Å 59.10Å 135.48Å 90.00° 91.12° 90.00°	Depositor
Resolution (Å)	41.21 – 2.50 41.21 – 2.50	Depositor EDS
% Data completeness (in resolution range)	98.3 (41.21-2.50) 98.8 (41.21-2.50)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.55 (at 2.51Å)	Xtriage
Refinement program	PHENIX (1.21.2_5419: ???)	Depositor
R, R_{free}	0.210 , 0.247 0.208 , 0.243	Depositor DCC
R_{free} test set	2359 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å^2)	50.0	Xtriage
Anisotropy	0.114	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.33 , 52.8	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.019 for h,-k,-l	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	8706	wwPDB-VP
Average B, all atoms (Å^2)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.33% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, EDO, ACT, CA, A1CZG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	A	0.12	0/4379	0.28	0/5954
1	B	0.10	0/4294	0.26	0/5829
All	All	0.11	0/8673	0.27	0/11783

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4295	0	4263	24	0
1	B	4217	0	4196	40	0
2	A	20	0	0	0	0
2	B	20	0	0	0	0
3	A	2	0	0	0	0
3	B	1	0	0	0	0
4	A	1	0	0	0	0
5	A	4	0	6	0	0
5	B	4	0	6	1	0
6	A	4	0	3	0	0
7	A	83	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	B	55	0	0	0	0
All	All	8706	0	8474	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:474:GLU:HG2	1:A:475:LEU:HD12	1.75	0.67
1:B:338:LYS:HD3	1:B:339:HIS:H	1.62	0.64
1:A:377:GLN:HB3	1:A:380:ARG:HG3	1.85	0.59
1:B:496:ASP:HB3	1:B:499:ALA:HB3	1.85	0.59
1:A:77:LEU:HD12	1:A:110:PRO:HG2	1.86	0.58
1:B:40:GLY:HA3	1:B:155:ARG:HG3	1.86	0.58
1:A:114:GLU:CD	1:A:114:GLU:H	2.15	0.55
1:A:22:VAL:HG21	1:A:149:ALA:HA	1.89	0.53
1:A:109:GLY:N	1:A:110:PRO:HD3	2.25	0.51
1:A:306:ILE:HD12	1:A:397:ILE:HG21	1.93	0.51
1:B:568:ALA:HA	1:B:572:LEU:HD22	1.93	0.50
1:A:77:LEU:HD11	1:A:112:HIS:CD2	2.46	0.50
1:B:442:LEU:HA	1:B:448:PRO:HA	1.94	0.50
1:B:107:LEU:HD12	1:B:110:PRO:HD3	1.94	0.49
1:B:556:ARG:HG2	1:B:563:THR:HG23	1.93	0.49
1:A:38:ALA:HB3	1:A:41:LEU:HD12	1.95	0.49
1:B:333:ASP:C	1:B:337:ALA:H	2.21	0.48
1:B:294:VAL:HG12	1:B:302:GLU:HB2	1.96	0.48
1:A:281:VAL:HG13	1:A:285:ASP:HB2	1.95	0.48
1:A:324:GLU:HG3	1:A:325:PRO:HA	1.97	0.47
1:B:77:LEU:HD21	1:B:115:LEU:HD22	1.96	0.47
1:B:509:ASP:O	1:B:513:LEU:HG	2.16	0.46
1:B:119:THR:O	1:B:123:GLU:HG2	2.17	0.45
1:B:197:VAL:HG12	1:B:232:ALA:HB1	1.99	0.45
1:B:543:VAL:HB	1:B:554:GLU:HB2	1.98	0.44
1:B:148:ARG:HH22	1:B:153:ARG:HH11	1.64	0.44
1:B:338:LYS:HD3	1:B:339:HIS:N	2.30	0.44
1:B:10:LEU:O	1:B:11:ARG:HD2	2.18	0.44
1:A:148:ARG:HH22	1:A:220:GLU:HG3	1.82	0.44
1:B:259:THR:HG22	1:B:271:TRP:CE2	2.53	0.44
1:B:2:ILE:HD11	1:B:507:ALA:C	2.43	0.44
1:A:112:HIS:O	1:A:116:PHE:HD1	2.00	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:112:HIS:HB2	1:A:116:PHE:HE1	1.82	0.43
1:B:546:ARG:H	5:B:603:EDO:H22	1.83	0.43
1:B:131:PRO:HB3	1:B:166:ASP:OD1	2.18	0.43
1:B:377:GLN:HB2	1:B:380:ARG:HG3	2.00	0.43
1:A:167:ILE:HD12	1:A:168:ASP:N	2.33	0.43
1:B:38:ALA:HB3	1:B:41:LEU:HD12	1.99	0.43
1:B:93:ASP:O	1:B:143:ASP:HB3	2.19	0.43
1:B:160:LYS:HE3	1:B:160:LYS:HB3	1.87	0.43
1:B:142:ARG:HD2	1:B:157:PHE:HZ	1.83	0.42
1:B:140:LYS:HD3	1:B:140:LYS:HA	1.90	0.42
1:B:214:ALA:HB2	1:B:423:ILE:HD13	2.00	0.42
1:B:338:LYS:CD	1:B:339:HIS:H	2.31	0.42
1:A:531:LYS:HE3	1:A:531:LYS:HB3	1.90	0.42
1:A:474:GLU:CD	1:A:474:GLU:H	2.28	0.42
1:A:14:ARG:HD2	1:A:14:ARG:HA	1.91	0.42
1:A:10:LEU:O	1:A:11:ARG:HD2	2.20	0.42
1:B:35:ARG:HD3	1:B:35:ARG:HA	1.93	0.42
1:B:234:ASN:OD1	1:B:236:GLU:HG3	2.20	0.41
1:A:543:VAL:HB	1:A:554:GLU:HB2	2.01	0.41
1:A:149:ALA:HB1	1:A:152:LEU:HD12	2.02	0.41
1:B:51:VAL:HG13	1:B:483:VAL:HG12	2.02	0.41
1:B:114:GLU:CD	1:B:114:GLU:H	2.29	0.41
1:B:527:GLY:O	1:B:531:LYS:HE2	2.21	0.41
1:B:303:LEU:HD11	1:B:331:LEU:HD11	2.03	0.41
1:B:541:ILE:HB	1:B:556:ARG:HB2	2.01	0.41
1:B:5:MET:HE2	1:B:5:MET:HB3	1.85	0.41
1:B:59:ILE:HG21	1:B:160:LYS:HG3	2.02	0.41
1:A:124:TYR:CD2	1:A:129:ASP:HB3	2.55	0.41
1:B:361:VAL:HB	1:B:365:ILE:HG13	2.01	0.40
1:B:15:ASP:HA	1:B:30:ARG:HD2	2.03	0.40
1:A:49:LEU:O	1:A:53:ARG:HG2	2.20	0.40
1:A:112:HIS:HB2	1:A:116:PHE:CE1	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	556/588 (95%)	537 (97%)	18 (3%)	1 (0%)	43 63
1	B	538/588 (92%)	514 (96%)	23 (4%)	1 (0%)	43 63
All	All	1094/1176 (93%)	1051 (96%)	41 (4%)	2 (0%)	43 63

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	110	PRO
1	A	207	SER

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	439/465 (94%)	436 (99%)	3 (1%)	76 89
1	B	432/465 (93%)	431 (100%)	1 (0%)	87 95
All	All	871/930 (94%)	867 (100%)	4 (0%)	81 92

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	128	LYS
1	A	167	ILE
1	A	516	GLU
1	B	281	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	296	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	425	HIS
1	A	472	HIS
1	B	428	GLN
1	B	562	GLN

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 9 ligands modelled in this entry, 4 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
6	ACT	A	606	-	3,3,3	1.14	0	3,3,3	1.24	0
5	EDO	B	603	-	3,3,3	0.25	0	2,2,2	0.35	0
5	EDO	A	605	-	3,3,3	0.25	0	2,2,2	0.24	0
2	A1CZG	A	601	-	21,21,21	0.16	0	25,29,29	0.37	0
2	A1CZG	B	601	-	21,21,21	0.20	0	25,29,29	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1CZG	B	601	-	-	2/12/12/12	0/2/2/2
5	EDO	B	603	-	-	1/1/1/1	-
5	EDO	A	605	-	-	1/1/1/1	-
2	A1CZG	A	601	-	-	2/12/12/12	0/2/2/2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	A1CZG	N6-C7-C9-C15
2	B	601	A1CZG	N6-C7-C9-C15
2	A	601	A1CZG	O8-C7-C9-C15
2	B	601	A1CZG	O8-C7-C9-C15
5	B	603	EDO	O1-C1-C2-O2
5	A	605	EDO	O1-C1-C2-O2


There are no ring outliers.

1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	603	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand A1CZG A 601




Bond lengths



Bond angles




Torsions




Rings


Ligand A1CZG B 601




Bond lengths



Bond angles



Torsions



Rings

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	564/588 (95%)	0.35	42 (7%)	20	18	30, 54, 108, 177	2 (0%)
1	B	552/588 (93%)	0.40	41 (7%)	20	18	31, 58, 114, 178	0
All	All	1116/1176 (94%)	0.38	83 (7%)	20	18	30, 56, 111, 178	2 (0%)

All (83) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	445	ASP	5.3
1	B	202	GLY	5.2
1	B	439	ALA	5.2
1	B	127	TYR	5.0
1	B	110	PRO	4.7
1	A	435	ASP	4.4
1	B	91	TYR	4.4
1	B	449	VAL	4.3
1	B	437	PHE	4.3
1	B	111	THR	4.2
1	A	110	PRO	4.0
1	A	433	TYR	3.9
1	B	124	TYR	3.9
1	A	276	ASP	3.6
1	A	432	LYS	3.6
1	B	443	GLY	3.5
1	A	15	ASP	3.3
1	A	109	GLY	3.3
1	A	436	ALA	3.3
1	B	567	VAL	3.2
1	B	167	ILE	3.1
1	A	330	LEU	3.1
1	A	334	ASP	3.1
1	A	22	VAL	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	582	GLY	3.0
1	B	438	THR	3.0
1	A	326	ALA	3.0
1	A	430	GLY	2.9
1	A	2	ILE	2.9
1	B	332	ASP	2.9
1	A	1	MET	2.9
1	A	169	ALA	2.8
1	A	447	LYS	2.8
1	A	201	SER	2.8
1	A	207	SER	2.8
1	B	448	PRO	2.7
1	A	80	ALA	2.7
1	B	430	GLY	2.7
1	A	431	SER	2.7
1	B	441	VAL	2.7
1	B	165	PHE	2.6
1	A	446	GLY	2.5
1	A	128	LYS	2.5
1	A	570	ALA	2.5
1	B	128	LYS	2.5
1	B	163	TYR	2.5
1	B	497	ALA	2.5
1	B	297	PRO	2.5
1	B	173	LYS	2.5
1	A	81	PRO	2.4
1	B	172	LEU	2.4
1	B	376	ASP	2.4
1	B	326	ALA	2.4
1	B	333	ASP	2.4
1	B	294	VAL	2.4
1	B	296	GLN	2.4
1	A	124	TYR	2.3
1	A	434	THR	2.3
1	A	437	PHE	2.3
1	B	130	PHE	2.3
1	A	449	VAL	2.3
1	A	111	THR	2.3
1	B	299	GLY	2.3
1	B	201	SER	2.3
1	A	452	THR	2.2
1	B	442	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	294	VAL	2.2
1	A	331	LEU	2.2
1	B	126	SER	2.2
1	A	17	PRO	2.2
1	A	325	PRO	2.2
1	B	570	ALA	2.2
1	A	450	ARG	2.2
1	B	107	LEU	2.1
1	B	295	ARG	2.1
1	B	410	ASP	2.1
1	B	227	LEU	2.1
1	A	442	LEU	2.1
1	A	327	ASP	2.1
1	B	229	SER	2.1
1	A	172	LEU	2.0
1	B	338	LYS	2.0
1	A	443	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
5	EDO	A	605	4/4	0.73	0.20	50,53,57,59	0
3	MG	A	603	1/1	0.81	0.21	78,78,78,78	0
5	EDO	B	603	4/4	0.83	0.19	74,75,80,80	0
2	A1CZG	B	601	20/20	0.85	0.10	53,65,86,101	0
2	A1CZG	A	601	20/20	0.85	0.11	44,57,79,93	0
4	CA	A	604	1/1	0.91	0.12	110,110,110,110	0

Continued on next page...


Continued from previous page...


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
6	ACT	A	606	4/4	0.93	0.14	40,44,50,53	0
3	MG	B	602	1/1	0.95	0.08	72,72,72,72	0
3	MG	A	602	1/1	0.96	0.08	52,52,52,52	1


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around A1CZG B 601:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)








Electron density around A1CZG A 601:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.