Full wwPDB X-ray Structure Validation Report ⓘ

Oct 25, 2026 – 11:24 AM EDT

PDB ID	:	9YYT / pdb_00009yyt
Title	:	Crystal structure of M. tuberculosis ProRS bound to compound 4c
Authors	:	Gade, P.; Michalska, K.; Joachimiak, A.; Center for Structural Biology of Infectious Diseases (CSBID)
Deposited on	:	2025-10-29
Resolution	:	2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references ⓘ) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.2
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1091 (2.54-2.54)
Clashscore	190562	1120 (2.54-2.54)
Ramachandran outliers	187476	1106 (2.54-2.54)
Sidechain outliers	187428	1106 (2.54-2.54)
RSRZ outliers	180081	1091 (2.54-2.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	588	6% 89% 10% .
1	B	588	6% 84% 12% .

2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 9065 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Proline–tRNA ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	578	Total	C	N	O	S	0	0	0
			4378	2763	772	835	8			
1	B	565	Total	C	N	O	S	0	1	0
			4340	2741	769	822	8			

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	583	GLU	-	expression tag	UNP P9WFT9
A	584	ASN	-	expression tag	UNP P9WFT9
A	585	LEU	-	expression tag	UNP P9WFT9
A	586	TYR	-	expression tag	UNP P9WFT9
A	587	PHE	-	expression tag	UNP P9WFT9
A	588	GLN	-	expression tag	UNP P9WFT9
B	583	GLU	-	expression tag	UNP P9WFT9
B	584	ASN	-	expression tag	UNP P9WFT9
B	585	LEU	-	expression tag	UNP P9WFT9
B	586	TYR	-	expression tag	UNP P9WFT9
B	587	PHE	-	expression tag	UNP P9WFT9
B	588	GLN	-	expression tag	UNP P9WFT9

- Molecule 2 is 3-(3-chlorobenzamido)pyrazine-2-carboxamide (CCD ID: A1CZL) (formula: $C_{12}H_9ClN_4O_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Cl	N	O	0	0
			19	12	1	4	2		
2	B	1	Total	C	Cl	N	O	0	0
			19	12	1	4	2		

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Mg 1 1	0	0
4	B	1	Total Mg 1 1	0	0

- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Ca 1 1	0	0
5	B	1	Total Ca 1 1	0	0

- Molecule 6 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0

- Molecule 7 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	154	Total O 154 154	0	0
7	B	131	Total O 131 131	0	0


3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Proline–tRNA ligase


Chain A:






• Molecule 1: Proline–tRNA ligase

Chain B:





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	86.05Å 59.75Å 137.66Å 90.00° 90.81° 90.00°	Depositor
Resolution (Å)	54.81 – 2.55 54.81 – 2.55	Depositor EDS
% Data completeness (in resolution range)	98.4 (54.81-2.55) 98.4 (54.81-2.55)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.55 (at 2.55Å)	Xtriage
Refinement program	PHENIX (1.21.2_5419: ???)	Depositor
R, R_{free}	0.230 , 0.271 0.230 , 0.273	Depositor DCC
R_{free} test set	2237 reflections (4.85%)	wwPDB-VP
Wilson B-factor (Å^2)	58.3	Xtriage
Anisotropy	0.361	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.27 , 34.7	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.014 for h,-k,-l	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	9065	wwPDB-VP
Average B, all atoms (Å^2)	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.76% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, MG, EDO, ACT, A1CZL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	A	0.09	0/4464	0.25	0/6074
1	B	0.12	0/4426	0.27	0/6016
All	All	0.11	0/8890	0.26	0/12090

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4378	0	4302	33	0
1	B	4340	0	4300	44	0
2	A	19	0	0	1	0
2	B	19	0	0	0	0
3	A	8	0	12	1	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	4	0	3	0	0
6	B	8	0	6	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	154	0	0	0	0
7	B	131	0	0	0	0
All	All	9065	0	8623	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:113:GLU:HG3	1:B:163:TYR:HE2	1.49	0.78
1:B:95:VAL:HG11	1:B:107:LEU:HD12	1.68	0.76
1:B:556:ARG:HG2	1:B:563:THR:HG22	1.72	0.69
1:A:377:GLN:HB3	1:A:380:ARG:HG3	1.79	0.64
1:B:375:ALA:HB3	1:B:380:ARG:HB3	1.82	0.61
1:A:324:GLU:HB2	1:A:325:PRO:HD2	1.83	0.60
1:B:312:ARG:HG2	1:B:403:ARG:HH22	1.66	0.59
1:B:494:ASN:ND2	1:B:546:ARG:H	1.99	0.59
1:B:490:LEU:HD23	1:B:519:LEU:HG	1.83	0.59
1:A:568:ALA:HA	1:A:572:LEU:HD22	1.85	0.58
1:A:543:VAL:HB	1:A:554:GLU:HB2	1.86	0.58
1:A:2:ILE:HD12	1:A:517:VAL:HB	1.87	0.57
1:A:446:GLY:HA3	1:B:313:GLU:HG2	1.88	0.55
1:B:490:LEU:HD12	1:B:542:VAL:HG23	1.91	0.53
1:B:8:LEU:HD21	1:B:47:LEU:HD21	1.91	0.53
1:B:116:PHE:O	1:B:120:VAL:HG23	2.09	0.52
1:A:288:LYS:HA	3:A:603:EDO:H12	1.91	0.52
1:B:496:ASP:HB3	1:B:499:ALA:HB3	1.91	0.51
1:A:312:ARG:HH12	1:A:403:ARG:HG2	1.76	0.51
1:B:519:LEU:CD2	1:B:521:ASP:HB3	2.42	0.49
1:B:180:ARG:O	1:B:184:GLN:HG3	2.13	0.49
1:B:113:GLU:HG3	1:B:163:TYR:CE2	2.38	0.49
1:B:296:GLN:HB3	1:B:297:PRO:HD2	1.95	0.48
1:A:59:ILE:HG21	1:A:160:LYS:HG3	1.95	0.48
1:B:2:ILE:HD12	1:B:517:VAL:HB	1.95	0.48
1:B:149:ALA:HB1	1:B:152:LEU:HD12	1.96	0.48
1:A:137:ILE:HG12	1:A:160:LYS:HG2	1.96	0.47
1:B:426:ILE:HG22	1:B:456:TYR:HD1	1.79	0.47
1:B:112:HIS:O	1:B:116:PHE:HD1	1.97	0.47
1:B:22:VAL:HG23	1:B:25:HIS:H	1.80	0.47
1:A:38:ALA:HB3	1:A:41:LEU:HD12	1.97	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:564:ARG:NH2	1:B:566:LEU:HD21	2.30	0.47
1:A:112:HIS:HB3	1:A:115:LEU:HB2	1.97	0.46
1:A:127:TYR:HB2	1:A:167:ILE:HG21	1.97	0.46
1:A:279:ARG:HD3	1:A:279:ARG:H	1.80	0.46
1:A:273:ASN:HD21	1:A:281:VAL:H	1.63	0.46
1:A:226:CYS:HB3	1:A:231:TYR:HB3	1.96	0.46
1:B:494:ASN:HD22	1:B:546:ARG:H	1.62	0.46
1:B:2:ILE:HD11	1:B:511:ASP:HB2	1.98	0.46
1:B:38:ALA:HB3	1:B:41:LEU:HB2	1.98	0.46
1:A:494:ASN:HB2	1:A:545:GLY:HA2	1.97	0.45
1:B:476:GLY:HA3	1:B:535:LEU:HA	1.97	0.45
1:B:240:THR:HB	1:B:363:PRO:HG2	1.99	0.45
1:A:154:ALA:H	2:A:601:A1CZL:C7	2.29	0.44
1:B:40:GLY:O	1:B:154:ALA:HB1	2.18	0.44
1:A:149:ALA:HB1	1:A:152:LEU:HD12	1.98	0.44
1:A:350:LYS:HB3	1:A:376:ASP:OD2	2.18	0.44
1:A:429:LEU:HD12	1:A:453:MET:HG3	2.00	0.44
1:B:197:VAL:HG12	1:B:232:ALA:HB1	2.00	0.44
1:A:49:LEU:O	1:A:53:ARG:HG2	2.18	0.43
1:B:471:HIS:O	1:B:478:ARG:HB2	2.18	0.43
1:B:429:LEU:HD12	1:B:453:MET:HG3	1.99	0.43
1:A:112:HIS:O	1:A:116:PHE:HD1	2.01	0.43
1:A:426:ILE:HG22	1:A:456:TYR:HD1	1.84	0.43
1:A:476:GLY:HA3	1:A:535:LEU:HA	2.01	0.43
1:B:246:LEU:HB2	1:B:368:GLY:HA3	2.00	0.42
1:B:25:HIS:O	1:B:29:ILE:HG12	2.19	0.42
1:B:71:ILE:HD11	1:B:116:PHE:CE2	2.55	0.42
1:B:259:THR:HG22	1:B:271:TRP:CE2	2.55	0.42
1:A:242:ARG:HG2	1:A:363:PRO:HB3	2.01	0.42
1:A:128:LYS:HE3	1:A:128:LYS:HB3	1.81	0.42
1:B:146:ARG:HG3	1:B:148:ARG:NH1	2.34	0.42
1:B:146:ARG:HG3	1:B:148:ARG:HH11	1.83	0.42
1:B:453:MET:HE2	1:B:453:MET:HB3	1.89	0.42
1:A:71:ILE:HD11	1:A:116:PHE:CE2	2.54	0.42
1:B:140:LYS:HD3	1:B:159:MET:HB2	2.01	0.42
1:A:331:LEU:HB3	1:A:335:ASP:HB2	2.02	0.41
1:A:77:LEU:HB2	1:A:107:LEU:HD12	2.03	0.41
1:A:471:HIS:O	1:A:478:ARG:HB2	2.21	0.41
1:B:245:THR:HG22	1:B:366:VAL:HB	2.03	0.41
1:B:478:ARG:HG2	1:B:558:ARG:HG3	2.03	0.41
1:A:324:GLU:O	1:A:325:PRO:C	2.64	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:442:LEU:HD11	1:B:446:GLY:HA2	2.02	0.41
1:B:503:ALA:HB2	1:B:544:VAL:HG11	2.02	0.41
1:A:292:ILE:HD12	1:A:328:TYR:HB2	2.02	0.40
1:B:138:GLN:HG2	1:B:139:THR:N	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	572/588 (97%)	549 (96%)	21 (4%)	2 (0%)	36	45
1	B	560/588 (95%)	533 (95%)	26 (5%)	1 (0%)	43	56
All	All	1132/1176 (96%)	1082 (96%)	47 (4%)	3 (0%)	36	45

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	325	PRO
1	A	432	LYS
1	B	546	ARG

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	443/465 (95%)	443 (100%)	0	100	100
1	B	445/465 (96%)	445 (100%)	0	100	100
All	All	888/930 (96%)	888 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	64	ASN
1	A	273	ASN
1	B	104	ASN
1	B	184	GLN
1	B	494	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 11 ligands modelled in this entry, 4 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\|>2$	Counts	RMSZ	$\#\|Z\|>2$
3	EDO	A	603	-	3,3,3	0.24	0	2,2,2	0.37	0
6	ACT	B	605	-	3,3,3	1.12	0	3,3,3	1.23	0
2	A1CZL	B	601	-	20,20,20	0.79	0	23,27,27	1.14	3 (13%)
6	ACT	B	604	-	3,3,3	1.12	0	3,3,3	1.21	0
2	A1CZL	A	601	-	20,20,20	0.80	1 (5%)	23,27,27	1.12	3 (13%)
3	EDO	A	602	-	3,3,3	0.25	0	2,2,2	0.35	0
6	ACT	A	606	-	3,3,3	1.14	0	3,3,3	1.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1CZL	B	601	-	-	0/12/12/12	0/2/2/2
3	EDO	A	603	-	-	0/1/1/1	-
2	A1CZL	A	601	-	-	0/12/12/12	0/2/2/2
3	EDO	A	602	-	-	0/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	A1CZL	C9-N8	2.02	1.38	1.34

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	601	A1CZL	C6-N5-C4	3.10	122.77	116.74
2	A	601	A1CZL	C6-N5-C4	3.05	122.67	116.74
2	B	601	A1CZL	C7-N8-C9	2.43	121.53	116.80
2	A	601	A1CZL	C7-N8-C9	2.36	121.38	116.80
2	B	601	A1CZL	C6-C7-N8	-2.12	119.20	122.19
2	A	601	A1CZL	C6-C7-N8	-2.00	119.37	122.19

There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.





2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	603	EDO	1	0
2	A	601	A1CZL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand A1CZL B 601



Ligand A1CZL A 601	
 Bond lengths	 Bond angles
 Torsions	 Rings

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	578/588 (98%)	0.45	36 (6%)	26	26	39, 63, 126, 163	0
1	B	565/588 (96%)	0.54	37 (6%)	25	24	35, 67, 128, 171	2 (0%)
All	All	1143/1176 (97%)	0.50	73 (6%)	25	24	35, 66, 127, 171	2 (0%)

All (73) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	91	TYR	4.7
1	A	110	PRO	4.6
1	A	325	PRO	4.0
1	B	430	GLY	3.9
1	A	431	SER	3.9
1	B	523	GLN	3.6
1	A	2	ILE	3.5
1	B	1	MET	3.4
1	A	109	GLY	3.4
1	B	110	PRO	3.4
1	B	127	TYR	3.4
1	B	449	VAL	3.3
1	A	433	TYR	3.3
1	A	276	ASP	3.2
1	A	202	GLY	3.2
1	B	444	GLU	3.1
1	A	443	GLY	3.0
1	B	91	TYR	3.0
1	B	124	TYR	3.0
1	B	555	LEU	2.9
1	B	442	LEU	2.9
1	A	1	MET	2.9
1	A	434	THR	2.9
1	A	586	TYR	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	356	ASN	2.8
1	A	584	ASN	2.8
1	B	441	VAL	2.8
1	B	570	ALA	2.7
1	A	430	GLY	2.6
1	A	203	ALA	2.6
1	B	507	ALA	2.6
1	A	107	LEU	2.5
1	A	585	LEU	2.5
1	A	297	PRO	2.4
1	B	563	THR	2.4
1	A	432	LYS	2.4
1	B	297	PRO	2.4
1	A	305	ALA	2.4
1	B	437	PHE	2.4
1	A	145	ALA	2.4
1	A	352	LEU	2.3
1	B	200	VAL	2.3
1	A	206	GLY	2.3
1	B	431	SER	2.3
1	B	443	GLY	2.3
1	B	425[A]	HIS	2.2
1	A	306	ILE	2.2
1	A	200	VAL	2.2
1	B	542	VAL	2.2
1	A	326	ALA	2.2
1	B	554	GLU	2.2
1	A	437	PHE	2.2
1	B	18	ALA	2.2
1	B	524	ALA	2.2
1	A	583	GLU	2.2
1	B	227	LEU	2.1
1	B	167	ILE	2.1
1	B	121	LYS	2.1
1	A	546	ARG	2.1
1	B	505	ALA	2.1
1	B	434	THR	2.1
1	B	448	PRO	2.1
1	B	577	ALA	2.1
1	B	548	TRP	2.1
1	B	92	GLY	2.1
1	A	83	GLU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	573	ALA	2.0
1	A	77	LEU	2.0
1	B	451	LEU	2.0
1	A	324	GLU	2.0
1	A	81	PRO	2.0
1	A	330	LEU	2.0
1	B	445	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
6	ACT	A	606	4/4	0.39	0.23	72,83,83,85	0
6	ACT	B	605	4/4	0.64	0.21	95,97,98,109	0
3	EDO	A	602	4/4	0.76	0.13	62,64,64,65	0
3	EDO	A	603	4/4	0.82	0.15	50,50,50,51	0
2	A1CZL	B	601	19/19	0.84	0.11	62,66,70,76	0
5	CA	B	603	1/1	0.88	0.08	89,89,89,89	0
2	A1CZL	A	601	19/19	0.89	0.10	55,59,63,66	0
5	CA	A	605	1/1	0.90	0.10	77,77,77,77	0
6	ACT	B	604	4/4	0.91	0.11	48,51,51,52	0
4	MG	B	602	1/1	0.92	0.16	74,74,74,74	0
4	MG	A	604	1/1	0.95	0.10	67,67,67,67	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around A1CZL B 601:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)










Electron density around A1CZL A 601:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







6.5 Other polymers (i)

There are no such residues in this entry.