Full wwPDB X-ray Structure Validation Report ⓘ

Oct 25, 2026 – 10:43 AM EDT

PDB ID	:	9YZX / pdb_00009yzx
Title	:	Crystal structure of M. tuberculosis ProRS bound to compound 4d
Authors	:	Gade, P.; Michalska, K.; Joachimiak, A.; Center for Structural Biology of Infectious Diseases (CSBID)
Deposited on	:	2025-10-30
Resolution	:	2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references ⓘ) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.2
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	588	2% 89% 10%
1	B	588	2% 87% 10%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PEG	A	603	-	-	X	-

2 Entry composition (i)

There are 9 unique types of molecules in this entry. The entry contains 9137 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Proline–tRNA ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	578	Total	C	N	O	S	0	0	0
			4403	2785	774	836	8			
1	B	576	Total	C	N	O	S	0	2	0
			4449	2808	789	844	8			

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	583	GLU	-	expression tag	UNP P9WFT9
A	584	ASN	-	expression tag	UNP P9WFT9
A	585	LEU	-	expression tag	UNP P9WFT9
A	586	TYR	-	expression tag	UNP P9WFT9
A	587	PHE	-	expression tag	UNP P9WFT9
A	588	GLN	-	expression tag	UNP P9WFT9
B	583	GLU	-	expression tag	UNP P9WFT9
B	584	ASN	-	expression tag	UNP P9WFT9
B	585	LEU	-	expression tag	UNP P9WFT9
B	586	TYR	-	expression tag	UNP P9WFT9
B	587	PHE	-	expression tag	UNP P9WFT9
B	588	GLN	-	expression tag	UNP P9WFT9

- Molecule 2 is 3-(4-fluorobenzamido)pyrazine-2-carboxamide (CCD ID: A1CZO) (formula: $C_{12}H_9FN_4O_2$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	0	0
			19	12	1	4	2		
2	B	1	Total	C	F	N	O	0	0
			19	12	1	4	2		

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O		0	0
			4	2	2			
3	B	1	Total	C	O		0	0
			4	2	2			

- Molecule 4 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 7 4 3	0	0


- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Mg 1 1	0	0
5	B	1	Total Mg 1 1	0	0

- Molecule 6 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Ca 1 1	0	0
6	B	1	Total Ca 1 1	0	0

- Molecule 7 is SULFATE ION (CCD ID: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total O S 5 4 1	0	0

- Molecule 8 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total C O 4 2 2	0	0

- Molecule 9 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	117	Total O 117 117	0	0
9	B	102	Total O 102 102	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Proline–tRNA ligase

Chain A:



• Molecule 1: Proline–tRNA ligase

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	86.16Å 59.67Å 137.83Å 90.00° 90.81° 90.00°	Depositor
Resolution (Å)	34.45 – 2.60 34.45 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.8 (34.45-2.60) 99.8 (34.45-2.60)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.76 (at 2.61Å)	Xtriage
Refinement program	PHENIX (1.21.2_5419: ???)	Depositor
R, R_{free}	0.226 , 0.269 0.226 , 0.269	Depositor DCC
R_{free} test set	2174 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å^2)	57.2	Xtriage
Anisotropy	0.146	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.30 , 38.0	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.014 for h,-k,-l	Xtriage
F_o,F_c correlation	0.93	EDS
Total number of atoms	9137	wwPDB-VP
Average B, all atoms (Å^2)	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.43% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, A1CZO, CA, SO4, PEG, ACT, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	A	0.12	0/4493	0.29	0/6116
1	B	0.14	0/4540	0.32	0/6175
All	All	0.13	0/9033	0.31	0/12291

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4403	0	4340	40	0
1	B	4449	0	4401	39	0
2	A	19	0	0	0	0
2	B	19	0	0	0	0
3	A	4	0	6	1	0
3	B	4	0	6	0	0
4	A	7	0	10	4	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	5	0	0	0	0
8	A	4	0	3	0	0
9	A	117	0	0	0	0
9	B	102	0	0	0	0
All	All	9137	0	8766	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:50:ARG:HG2	4:A:603:PEG:H31	1.64	0.80
1:B:503:ALA:HB2	1:B:544:VAL:HG21	1.64	0.78
1:A:354:GLU:HA	1:B:546:ARG:HH22	1.50	0.75
1:A:429:LEU:HD11	1:A:453:MET:HG3	1.71	0.73
1:B:490:LEU:HD12	1:B:542:VAL:HB	1.78	0.64
1:A:38:ALA:HB3	1:A:41:LEU:HD12	1.84	0.60
1:A:113:GLU:HG3	1:A:163:TYR:HE2	1.66	0.59
1:A:191:GLN:HG3	1:A:480:PRO:HB3	1.84	0.58
1:B:434:THR:HG22	1:B:439:ALA:HB3	1.84	0.58
1:B:550:ASP:HB3	1:B:552:VAL:HG22	1.85	0.57
1:A:272:ALA:HB3	1:A:281:VAL:HG21	1.87	0.56
1:B:125:SER:C	1:B:441:VAL:HG13	2.31	0.56
1:B:474:GLU:H	1:B:474:GLU:CD	2.14	0.55
1:B:113:GLU:HG3	1:B:163:TYR:HE2	1.71	0.54
1:B:38:ALA:HB3	1:B:41:LEU:HD12	1.89	0.54
1:A:53:ARG:HB2	4:A:603:PEG:H42	1.90	0.54
1:A:494:ASN:HB2	1:A:545:GLY:HA2	1.89	0.54
1:B:435:ASP:HA	1:B:450:ARG:HH11	1.73	0.54
1:B:441:VAL:HG12	1:B:442:LEU:N	2.23	0.53
1:A:568:ALA:HA	1:A:572:LEU:HD22	1.91	0.52
1:A:503:ALA:HB2	1:A:544:VAL:HG11	1.90	0.52
1:B:2:ILE:HD12	1:B:517:VAL:HB	1.90	0.52
1:B:490:LEU:HD23	1:B:519:LEU:HD23	1.91	0.52
1:B:295:ARG:HH22	1:B:297:PRO:HA	1.75	0.52
1:A:377:GLN:HB2	1:A:380:ARG:HG3	1.94	0.50
1:A:112:HIS:CE1	1:A:140:LYS:HE3	2.47	0.50
1:B:441:VAL:O	1:B:442:LEU:HB2	2.11	0.49
1:B:435:ASP:HA	1:B:450:ARG:NH1	2.28	0.49
1:B:102:ARG:HB3	1:B:104:ASN:OD1	2.13	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:113:GLU:HG3	1:A:163:TYR:CE2	2.47	0.48
1:B:51:VAL:HG13	1:B:483:VAL:HG12	1.95	0.48
1:B:490:LEU:HB3	1:B:519:LEU:HD23	1.95	0.48
1:B:425[B]:HIS:NE2	1:B:457:GLY:HA3	2.29	0.47
1:A:53:ARG:HG3	4:A:603:PEG:H21	1.95	0.47
1:B:160:LYS:O	1:B:457:GLY:HA2	2.15	0.47
1:B:116:PHE:O	1:B:120:VAL:HG23	2.15	0.47
1:A:312:ARG:HH12	1:A:403:ARG:HG2	1.79	0.46
1:A:5:MET:HE2	1:A:5:MET:HB3	1.79	0.46
1:A:259:THR:HG22	1:A:271:TRP:CZ2	2.51	0.46
1:B:425[B]:HIS:ND1	1:B:427:PHE:HE1	2.14	0.46
1:A:374:GLY:N	3:A:602:EDO:H21	2.31	0.45
1:B:541:ILE:HB	1:B:556:ARG:HB2	1.98	0.45
1:B:259:THR:HG22	1:B:271:TRP:CE2	2.52	0.45
1:B:493:ALA:HB1	1:B:526:PRO:HB3	1.99	0.45
1:A:226:CYS:HB3	1:A:231:TYR:HB3	1.98	0.44
1:B:292:ILE:HG22	1:B:330:LEU:HD22	1.99	0.44
1:A:39:PRO:O	1:A:155:ARG:HD3	2.17	0.44
1:B:104:ASN:HB2	1:B:106:TYR:CE2	2.52	0.44
1:A:56:GLU:HG2	1:A:460:VAL:HG11	2.00	0.43
1:B:295:ARG:NH2	1:B:297:PRO:HA	2.32	0.43
1:B:361:VAL:HB	1:B:365:ILE:HG13	1.99	0.43
1:A:552:VAL:HA	1:A:566:LEU:O	2.19	0.43
1:A:50:ARG:HA	4:A:603:PEG:H22	2.00	0.43
1:A:417:MET:HE3	1:A:417:MET:HB3	1.85	0.43
1:A:87:ARG:HA	1:A:90:GLN:HB3	2.01	0.42
1:A:496:ASP:OD2	1:A:498:GLN:HG2	2.19	0.42
1:B:306:ILE:HA	1:B:360:LEU:O	2.20	0.42
1:A:293:LYS:HG2	1:A:331:LEU:HD12	2.01	0.42
1:B:510:LEU:HA	1:B:513:LEU:HD23	2.01	0.42
1:A:152:LEU:HD23	1:A:152:LEU:HA	1.92	0.42
1:B:224:VAL:HB	1:B:415:LEU:HD22	2.01	0.42
1:B:426:ILE:HG22	1:B:456:TYR:HD1	1.84	0.42
1:A:272:ALA:CB	1:A:281:VAL:HG21	2.48	0.42
1:B:292:ILE:HD11	1:B:304:LEU:HD23	2.01	0.42
1:A:404:GLU:HG2	1:A:417:MET:CG	2.50	0.42
1:A:38:ALA:HB1	1:A:39:PRO:HD2	2.02	0.41
1:A:259:THR:HG22	1:A:271:TRP:CE2	2.55	0.41
1:A:404:GLU:HG2	1:A:417:MET:HG3	2.02	0.41
1:B:88:TRP:CZ3	1:B:107:LEU:HB3	2.55	0.41
1:A:167:ILE:H	1:A:167:ILE:HG13	1.70	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:261:ASP:OD1	1:A:263:PRO:HD3	2.20	0.41
1:A:5:MET:HE3	1:A:10:LEU:HD23	2.02	0.41
1:A:543:VAL:HB	1:A:554:GLU:HB2	2.02	0.41
1:B:128:LYS:H	1:B:128:LYS:HG3	1.69	0.41
1:B:84:THR:HG22	1:B:436:ALA:HB1	2.03	0.40
1:A:159:MET:HE3	1:A:159:MET:HB3	1.83	0.40
1:B:69:GLN:HB2	1:B:134:LEU:HD23	2.03	0.40
1:A:471:HIS:O	1:A:478:ARG:HB2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	574/588 (98%)	567 (99%)	7 (1%)	0	100	100
1	B	574/588 (98%)	562 (98%)	11 (2%)	1 (0%)	43	66
All	All	1148/1176 (98%)	1129 (98%)	18 (2%)	1 (0%)	48	70

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	442	LEU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	447/465 (96%)	445 (100%)	2 (0%)	84 93
1	B	457/465 (98%)	454 (99%)	3 (1%)	76 89
All	All	904/930 (97%)	899 (99%)	5 (1%)	78 91

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	89	THR
1	A	438	THR
1	B	519	LEU
1	B	553	VAL
1	B	572	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	25	HIS
1	A	339	HIS
1	A	562	GLN
1	B	69	GLN
1	B	191	GLN
1	B	523	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 11 ligands modelled in this entry, 4 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	EDO	B	602	-	3,3,3	0.23	0	2,2,2	0.50	0
3	EDO	A	602	-	3,3,3	0.22	0	2,2,2	0.48	0
7	SO4	A	606	-	4,4,4	0.69	0	6,6,6	0.10	0
4	PEG	A	603	-	6,6,6	0.26	0	5,5,5	0.19	0
8	ACT	A	607	-	3,3,3	1.13	0	3,3,3	1.25	0
2	A1CZO	A	601	-	20,20,20	0.17	0	23,27,27	0.36	0
2	A1CZO	B	601	-	20,20,20	0.16	0	23,27,27	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	B	602	-	-	1/1/1/1	-
3	EDO	A	602	-	-	1/1/1/1	-
4	PEG	A	603	-	-	4/4/4/4	-
2	A1CZO	A	601	-	-	1/12/12/12	0/2/2/2
2	A1CZO	B	601	-	-	0/12/12/12	0/2/2/2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	603	PEG	O1-C1-C2-O2
4	A	603	PEG	O2-C3-C4-O4
3	A	602	EDO	O1-C1-C2-O2
2	A	601	A1CZO	C9-C4-N3-C2
4	A	603	PEG	C4-C3-O2-C2
3	B	602	EDO	O1-C1-C2-O2
4	A	603	PEG	C1-C2-O2-C3


There are no ring outliers.

2 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	602	EDO	1	0
4	A	603	PEG	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand A1CZO A 601




Bond lengths



Bond angles




Torsions




Rings


Ligand A1CZO B 601




Bond lengths



Bond angles



Torsions



Rings

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	578/588 (98%)	0.24	14 (2%)	59	54	36, 58, 102, 135	0
1	B	576/588 (97%)	0.31	13 (2%)	61	55	31, 64, 115, 136	2 (0%)
All	All	1154/1176 (98%)	0.28	27 (2%)	61	55	31, 61, 110, 136	2 (0%)

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	430	GLY	4.1
1	A	200	VAL	3.5
1	B	142	ARG	3.4
1	B	201	SER	3.2
1	B	90	GLN	3.0
1	A	248	ILE	2.9
1	B	275	ALA	2.7
1	B	208	ALA	2.6
1	A	275	ALA	2.5
1	A	249	ASP	2.4
1	B	493	ALA	2.4
1	A	433	TYR	2.4
1	A	555	LEU	2.4
1	A	277	LEU	2.3
1	B	84	THR	2.3
1	A	296	GLN	2.3
1	B	227	LEU	2.3
1	A	199	ALA	2.3
1	A	201	SER	2.3
1	B	89	THR	2.2
1	A	300	ASP	2.2
1	B	91	TYR	2.2
1	B	442	LEU	2.1
1	A	202	GLY	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	1	MET	2.0
1	B	449	VAL	2.0
1	A	297	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
8	ACT	A	607	4/4	0.65	0.16	56,56,58,60	0
7	SO4	A	606	5/5	0.82	0.14	92,93,95,97	0
2	A1CZO	B	601	19/19	0.85	0.10	49,52,56,59	0
4	PEG	A	603	7/7	0.86	0.18	40,40,43,44	0
2	A1CZO	A	601	19/19	0.88	0.09	51,54,56,60	0
3	EDO	A	602	4/4	0.90	0.11	42,43,43,44	0
3	EDO	B	602	4/4	0.91	0.11	45,45,46,49	0
5	MG	A	604	1/1	0.93	0.13	70,70,70,70	0
6	CA	A	605	1/1	0.94	0.07	62,62,62,62	0
6	CA	B	604	1/1	0.95	0.10	64,64,64,64	0
5	MG	B	603	1/1	0.97	0.12	65,65,65,65	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around A1CZO B 601:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)










Electron density around A1CZO A 601:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







6.5 Other polymers (i)

There are no such residues in this entry.