



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 30, 2026 – 06:08 PM EDT

PDB ID : 9ZGA / pdb\_00009zga  
Title : Crystal structure of DH511.12P Fab in complex with HIV-1 gp41 MPER peptide and phosphatidic acid (06:0 PA); double-lipid form  
Authors : Cho, S.Y.; Wilson, I.A.  
Deposited on : 2025-12-02  
Resolution : 1.97 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.015 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.50


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1506 (1.98-1.98)
Clashscore	190562	1534 (1.98-1.98)
Ramachandran outliers	187476	1518 (1.98-1.98)
Sidechain outliers	187428	1518 (1.98-1.98)
RSRZ outliers	180081	1506 (1.98-1.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	236	 5% 93% 6%
2	B	217	 2% 89% 11%
3	C	19	 16% 95% 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	PEG	B	302	-	-	X	-

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 7448 atoms, of which 3423 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Antibody DH511.12P Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	234	3438	1119	1682	292	338	7	0	0	0

- Molecule 2 is a protein called Antibody DH511.12P Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
2	B	217	3243	1036	1586	278	337	6	0	0	0

- Molecule 3 is a protein called Transmembrane protein gp41.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
3	C	19	329	120	155	29	25	0	0	0

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	668	LYS	-	expression tag	UNP Q73372
C	669	LYS	-	expression tag	UNP Q73372
C	670	LYS	-	expression tag	UNP Q73372
C	684	LYS	-	expression tag	UNP Q73372
C	685	LYS	-	expression tag	UNP Q73372
C	686	LYS	-	expression tag	UNP Q73372

- Molecule 4 is PHOSPHATE ION (CCD ID: PO4) (formula: O<sub>4</sub>P) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 5 is (2R)-3-(phosphonoxy)propane-1,2-diyl dihexanoate (CCD ID: 44E) (formula:  $C_{15}H_{29}O_8P$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	O	P	0	0
			18	10	7	1		
5	B	1	Total	C	O	P	0	0
			18	9	8	1		

- Molecule 6 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula:  $C_4H_{10}O_3$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 7 4 3	0	0
6	B	1	Total C O 7 4 3	0	0
6	C	1	Total C O 7 4 3	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	214	Total O 214 214	0	0
7	B	158	Total O 158 158	0	0
7	C	4	Total O 4 4	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Antibody DH511.12P Fab heavy chain



- Molecule 2: Antibody DH511.12P Fab light chain



- Molecule 3: Transmembrane protein gp41



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.53Å 73.53Å 230.41Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	32.88 – 1.97 32.88 – 1.97	Depositor EDS
% Data completeness (in resolution range)	99.6 (32.88-1.97) 99.6 (32.88-1.97)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.68 (at 1.97Å)	Xtrriage
Refinement program	PHENIX 1.21.2_5419	Depositor
R, $R_{free}$	0.226 , 0.264 0.225 , 0.262	Depositor DCC
$R_{free}$ test set	2488 reflections (4.74%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	29.0	Xtrriage
Anisotropy	0.221	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 35.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.031 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	7448	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.16% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, 44E, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.47	0/1804	0.60	0/2467
2	B	0.50	0/1691	0.66	0/2294
3	C	0.50	0/181	0.67	0/245
All	All	0.49	0/3676	0.63	0/5006

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1756	1682	1697	11	0
2	B	1657	1586	1596	16	0
3	C	174	155	160	1	0
4	A	5	0	0	0	0
5	A	18	0	18	1	0
5	B	18	0	13	2	0
6	A	7	0	10	2	0
6	B	7	0	10	4	0
6	C	7	0	10	0	0
7	A	214	0	0	1	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	B	158	0	0	0	0
7	C	4	0	0	0	0
All	All	4025	3423	3514	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:81:HIS:HB3	6:A:303:PEG:H12	1.74	0.68
1:A:12:VAL:HG11	1:A:82(C):LEU:HD13	1.82	0.60
2:B:110:VAL:HG21	6:B:302:PEG:H32	1.85	0.57
1:A:123:PRO:HD3	1:A:209:LYS:HE2	1.86	0.56
2:B:95(B):MET:HE2	2:B:97:ILE:HD11	1.88	0.56
2:B:32:TYR:CE2	5:B:301:44E:H4	2.41	0.56
1:A:100(H):TYR:CD2	5:B:301:44E:H5	2.42	0.54
1:A:143:LYS:HE3	7:A:408:HOH:O	2.06	0.54
2:B:110:VAL:CG2	6:B:302:PEG:H32	2.39	0.53
2:B:95:VAL:O	2:B:95(B):MET:HG3	2.10	0.51
2:B:39:ARG:HG2	2:B:84:ALA:HB2	1.93	0.50
2:B:161:GLU:HG2	2:B:175:LEU:HD21	1.95	0.49
2:B:33:LEU:HG	2:B:34:ASN:N	2.31	0.46
2:B:33:LEU:HD22	2:B:71:PHE:CG	2.51	0.45
1:A:9:GLY:HA2	1:A:18:LEU:HD21	1.98	0.45
5:A:302:44E:H28	3:C:680:TRP:CD1	2.51	0.45
2:B:124:GLN:HG2	2:B:129:THR:O	2.17	0.45
2:B:19:ILE:HD11	2:B:75:ILE:HD12	1.99	0.45
2:B:-2:SER:HB3	2:B:0:THR:H	1.81	0.45
2:B:110:VAL:HG21	6:B:302:PEG:H11	1.97	0.45
1:A:143:LYS:NZ	1:A:171:GLN:OE1	2.51	0.44
1:A:178:LEU:C	1:A:178:LEU:HD12	2.42	0.43
2:B:37:LYS:HE3	2:B:45:LYS:HD3	1.99	0.43
1:A:82:MET:CE	1:A:90:TYR:CZ	3.01	0.43
1:A:81:HIS:CG	6:A:303:PEG:H41	2.54	0.43
2:B:125:LEU:O	2:B:183:LYS:HD2	2.19	0.42
2:B:140:TYR:CG	2:B:141:PRO:HA	2.55	0.42
6:B:302:PEG:H32	6:B:302:PEG:H11	1.63	0.41
1:A:82:MET:HE1	1:A:90:TYR:CE1	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	232/236 (98%)	227 (98%)	5 (2%)	0	100	100
2	B	215/217 (99%)	206 (96%)	9 (4%)	0	100	100
3	C	17/19 (90%)	16 (94%)	1 (6%)	0	100	100
All	All	464/472 (98%)	449 (97%)	15 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	193/201 (96%)	191 (99%)	2 (1%)	68	65
2	B	186/192 (97%)	186 (100%)	0	100	100
3	C	15/19 (79%)	15 (100%)	0	100	100
All	All	394/412 (96%)	392 (100%)	2 (0%)	81	79

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	178	LEU
1	A	191	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	PEG	A	303	-	6,6,6	0.28	0	5,5,5	0.39	0
6	PEG	B	302	-	6,6,6	0.27	0	5,5,5	0.42	0
4	PO4	A	301	-	4,4,4	1.53	1 (25%)	6,6,6	1.00	0
5	44E	B	301	-	17,17,23	1.85	3 (17%)	18,21,28	1.63	2 (11%)
6	PEG	C	701	-	6,6,6	0.32	0	5,5,5	0.21	0
5	44E	A	302	-	17,17,23	1.21	2 (11%)	19,21,28	1.27	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PEG	A	303	-	-	3/4/4/4	-
6	PEG	B	302	-	-	3/4/4/4	-
5	44E	B	301	-	-	7/18/18/25	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PEG	C	701	-	-	4/4/4/4	-
5	44E	A	302	-	-	8/18/18/25	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	301	44E	O21-C2	-5.38	1.39	1.46
5	B	301	44E	O21-C21	3.38	1.40	1.33
5	A	302	44E	O21-C2	-2.74	1.40	1.46
5	B	301	44E	O31-C31	2.61	1.41	1.33
5	A	302	44E	O21-C21	2.47	1.41	1.34
4	A	301	PO4	P-O1	2.31	1.56	1.50

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	301	44E	O21-C21-O22	-5.27	118.94	125.70
5	A	302	44E	O21-C21-C22	4.18	120.52	111.48
5	B	301	44E	O11-P-O12	2.34	112.77	106.44

There are no chirality outliers.

All (25) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	301	44E	C1-O11-P-O13
5	B	301	44E	C1-O11-P-O14
5	B	301	44E	O22-C21-O21-C2
5	A	302	44E	C21-C22-C23-C4
6	B	302	PEG	O1-C1-C2-O2
6	B	302	PEG	C1-C2-O2-C3
6	C	701	PEG	O1-C1-C2-O2
5	A	302	44E	C23-C4-C5-C6
6	A	303	PEG	O2-C3-C4-O4
5	B	301	44E	C2-C1-O11-P
6	B	302	PEG	O2-C3-C4-O4
5	B	301	44E	O11-C1-C2-C3
5	B	301	44E	C1-O11-P-O12
5	A	302	44E	C1-C2-C3-O31
5	B	301	44E	O11-C1-C2-O21
5	A	302	44E	C2-C3-O31-C31
6	A	303	PEG	C4-C3-O2-C2

*Continued on next page...*

*Continued from previous page...*



Mol	Chain	Res	Type	Atoms
6	C	701	PEG	C4-C3-O2-C2
6	A	303	PEG	O1-C1-C2-O2
5	A	302	44E	O21-C2-C3-O31
5	A	302	44E	C1-O11-P-O13
6	C	701	PEG	O2-C3-C4-O4
6	C	701	PEG	C1-C2-O2-C3
5	A	302	44E	O21-C21-C22-C23
5	A	302	44E	O22-C21-C22-C23


There are no ring outliers.



4 monomers are involved in 9 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	303	PEG	2	0
6	B	302	PEG	4	0
5	B	301	44E	2	0
5	A	302	44E	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	234/236 (99%)	-0.07	12 (5%) 33 43	19, 29, 61, 86	0
2	B	217/217 (100%)	0.01	4 (1%) 67 76	22, 35, 52, 86	0
3	C	19/19 (100%)	0.74	3 (15%) 5 6	24, 36, 75, 79	0
All	All	470/472 (99%)	0.00	19 (4%) 42 52	19, 32, 56, 86	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	133	GLY	7.8
1	A	215	SER	5.3
1	A	128	SER	4.7
3	C	668	LYS	4.5
1	A	135	THR	4.2
1	A	130	SER	4.0
1	A	129	LYS	3.7
1	A	131	THR	3.7
2	B	212	CYS	3.3
1	A	134	GLY	3.2
2	B	-1	VAL	3.0
1	A	1	GLN	2.9
3	C	685	LYS	2.9
3	C	686	LYS	2.9
1	A	132	SER	2.8
2	B	-2	SER	2.7
1	A	127	SER	2.7
2	B	78	LEU	2.3
1	A	214	LYS	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	44E	A	302	18/24	0.72	0.26	40,58,92,101	0
5	44E	B	301	18/24	0.78	0.20	40,50,71,83	0
6	PEG	B	302	7/7	0.78	0.30	56,73,85,87	0
6	PEG	A	303	7/7	0.85	0.13	32,39,47,59	0
6	PEG	C	701	7/7	0.87	0.13	33,38,44,45	0
4	PO4	A	301	5/5	0.91	0.12	47,51,57,106	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around 44E A 302:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around 44E B 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







**Electron density around PEG B 302:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around PEG A 303:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.