Full wwPDB EM Validation Report ⓘ

Oct 25, 2026 – 11:32 AM EDT

PDB ID	:	9ZUD / pdb_00009zud
EMDB ID	:	EMD-74807
Title	:	EBV gB trimer with "deD" mutations (intermediate-1 conformation)
Authors	:	McCallum, M.; Veesler, D.; Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on	:	2025-12-23
Resolution	:	4.10 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references ⓘ) were used in the production of this report:


EMDB validation analysis	:	0.0.1.dev133
Mogul	:	2022.3.0, CSD as543be (2022)
MolProbity	:	4-5-2 with Phenix2.2
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics	:	202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 4.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	6458 (3.60 - 4.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	648	26% 64% · 32%
1	B	648	29% 45% · 53%
1	C	648	14% 50% · 48%

2 Entry composition


There are 2 unique types of molecules in this entry. The entry contains 6572 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called EBV gB trimer with "deD" mutations (intermediate-1 conformation).

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	438	Total	C	N	O	S	0	0
			2669	1676	500	477	16		
1	B	306	Total	C	N	O	S	0	0
			1650	999	321	318	12		
1	C	337	Total	C	N	O	S	0	0
			2057	1298	383	362	14		

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				AltConf
2	A	1	Total	C	N	O	0
			14	8	1	5	
2	A	1	Total	C	N	O	0
			14	8	1	5	
2	A	1	Total	C	N	O	0
			14	8	1	5	

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				AltConf
2	A	1	Total	C	N	O	0
			14	8	1	5	
2	A	1	Total	C	N	O	0
			14	8	1	5	
2	A	1	Total	C	N	O	0
			14	8	1	5	
2	B	1	Total	C	N	O	0
			14	8	1	5	
2	B	1	Total	C	N	O	0
			14	8	1	5	
2	B	1	Total	C	N	O	0
			14	8	1	5	
2	C	1	Total	C	N	O	0
			14	8	1	5	
2	C	1	Total	C	N	O	0
			14	8	1	5	
2	C	1	Total	C	N	O	0
			14	8	1	5	
2	C	1	Total	C	N	O	0
			14	8	1	5	
2	C	1	Total	C	N	O	0
			14	8	1	5	

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: EBV gB trimer with "deD" mutations (intermediate-1 conformation)


Chain A:



• Molecule 1: EBV gB trimer with "deD" mutations (intermediate-1 conformation)

Chain B:







• Molecule 1: EBV gB trimer with "deD" mutations (intermediate-1 conformation)

Chain C:






THR ASN TYR LYS ILE VAL ASP TYR ASP ASN ARG GLY THR ASN PRO GLN GLY GLU ARG ARG ALA PHE LEU ASP LYS GLY THR TYR THR LEU SER TRP LYS LEU GLU ASN ARG THR ALA
Y294 C295 P296 K313 S314 D320 GLU GLY T323 S324 S325 F326 I334 E335 L336 P337 D338 A339
G365 G377 L378 W382 L383 P384 L385 T386 L399 THR THR LEU ASN ASN PRO A457 R495 E496 I500 N501 S508 I509 Y510 G511 K512 A515 A516 K517 I523 S524 V525 S526 C527 C528 V529 M542 R543 V544 P545 G546 S547 E548 T549 R554 P555 L556 V557 N563
D564 T565 T581 K582 K583 M584 T585 E586 V587 A590 F596 Y605 D617 L622 Q623 T624 F625 I626 S627 L628 N629 T630 S631 L632 I633 C634 N635 I636 D637 F638 A639 S640 L641 E642 L643 Y644 SER ARG ASP GLU GLN ARG ALA SER ASN VAL PHE ASP LEU GLU GLY ILE PHE ARG
GLU TYR ASN PHE GLN ALA GLN ASN ILE ALA GLY LEU ARG LYS ASP LEU ASP ASN ALA VAL SER GLY SER GLY SER GLY GLY SER GLY HIS HIS HIS HIS HIS HIS HIS

4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	324509	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/Å^2$)	50	Depositor
Minimum defocus (nm)	400	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	1.005	Depositor
Minimum map value	-0.264	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.020	Depositor
Recommended contour level	0.3	Depositor
Map size (Å)	337.2, 337.2, 337.2	wwPDB
Map dimensions	400, 400, 400	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.84300005, 0.84300005, 0.84300005	Depositor

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|>5$	RMSZ	#$\|Z\|>5$
1	A	0.48	0/2713	0.93	28/3730 (0.8%)
1	B	0.50	0/1661	0.98	18/2299 (0.8%)
1	C	0.50	0/2096	0.90	18/2887 (0.6%)
All	All	0.49	0/6470	0.93	64/8916 (0.7%)

There are no bond length outliers.

All (64) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	336	LEU	CA-C-N	7.50	127.41	120.21
1	B	336	LEU	C-N-CA	7.50	127.41	120.21
1	C	529	VAL	CA-C-N	7.45	127.45	119.78
1	C	529	VAL	C-N-CA	7.45	127.45	119.78
1	A	169	LYS	CA-C-N	7.24	127.16	119.85
1	A	169	LYS	C-N-CA	7.24	127.16	119.85
1	A	501	ASN	CA-C-N	7.09	127.40	119.32
1	A	501	ASN	C-N-CA	7.09	127.40	119.32
1	C	68	CYS	CA-C-N	7.07	127.06	119.78
1	C	68	CYS	C-N-CA	7.07	127.06	119.78
1	B	46	PHE	CA-C-N	7.06	127.51	120.52
1	B	46	PHE	C-N-CA	7.06	127.51	120.52
1	B	383	LEU	CA-C-N	7.01	126.77	119.76
1	B	383	LEU	C-N-CA	7.01	126.77	119.76
1	C	46	PHE	CA-C-N	6.95	127.12	120.31
1	C	46	PHE	C-N-CA	6.95	127.12	120.31
1	A	46	PHE	CA-C-N	6.94	127.39	120.52
1	A	46	PHE	C-N-CA	6.94	127.39	120.52
1	B	529	VAL	CA-C-N	6.90	126.88	119.78
1	B	529	VAL	C-N-CA	6.90	126.88	119.78
1	A	383	LEU	CA-C-N	6.83	126.75	119.85

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	383	LEU	C-N-CA	6.83	126.75	119.85
1	A	529	VAL	CA-C-N	6.73	126.65	119.85
1	A	529	VAL	C-N-CA	6.73	126.65	119.85
1	A	455	ASN	CA-C-N	6.70	126.39	119.82
1	A	455	ASN	C-N-CA	6.70	126.39	119.82
1	A	336	LEU	CA-C-N	6.69	126.67	119.78
1	A	336	LEU	C-N-CA	6.69	126.67	119.78
1	C	554	ARG	CA-C-N	6.69	126.67	119.78
1	C	554	ARG	C-N-CA	6.69	126.67	119.78
1	A	233	SER	CA-C-N	6.61	126.75	119.87
1	A	233	SER	C-N-CA	6.61	126.75	119.87
1	A	68	CYS	CA-C-N	6.55	126.52	120.03
1	A	68	CYS	C-N-CA	6.55	126.52	120.03
1	B	501	ASN	CA-C-N	6.55	126.79	119.32
1	B	501	ASN	C-N-CA	6.55	126.79	119.32
1	B	554	ARG	CA-C-N	6.45	126.48	119.90
1	B	554	ARG	C-N-CA	6.45	126.48	119.90
1	A	554	ARG	CA-C-N	6.40	126.37	119.78
1	A	554	ARG	C-N-CA	6.40	126.37	119.78
1	A	295	CYS	CA-C-N	6.16	126.05	119.28
1	A	295	CYS	C-N-CA	6.16	126.05	119.28
1	C	501	ASN	CA-C-N	6.11	126.22	119.87
1	C	501	ASN	C-N-CA	6.11	126.22	119.87
1	A	90	ILE	CA-C-N	6.00	126.02	119.90
1	A	90	ILE	C-N-CA	6.00	126.02	119.90
1	B	386	THR	CA-C-N	5.92	125.94	119.90
1	B	386	THR	C-N-CA	5.92	125.94	119.90
1	C	336	LEU	CA-C-N	5.82	127.21	120.98
1	C	336	LEU	C-N-CA	5.82	127.21	120.98
1	C	386	THR	CA-C-N	5.79	125.81	119.90
1	C	386	THR	C-N-CA	5.79	125.81	119.90
1	A	386	THR	CA-C-N	5.76	126.10	119.93
1	A	386	THR	C-N-CA	5.76	126.10	119.93
1	B	544	VAL	CA-C-N	5.50	125.81	119.93
1	B	544	VAL	C-N-CA	5.50	125.81	119.93
1	C	295	CYS	CA-C-N	5.43	126.62	119.84
1	C	295	CYS	C-N-CA	5.43	126.62	119.84
1	B	295	CYS	CA-C-N	5.42	126.62	119.84
1	B	295	CYS	C-N-CA	5.42	126.62	119.84
1	A	544	VAL	CA-C-N	5.39	125.56	119.90
1	A	544	VAL	C-N-CA	5.39	125.56	119.90
1	C	544	VAL	CA-C-N	5.19	125.43	119.83

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	C	544	VAL	C-N-CA	5.19	125.43	119.83

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2669	0	1796	4	0
1	B	1650	0	902	3	0
1	C	2057	0	1373	4	0
2	A	84	0	78	0	0
2	B	42	0	39	2	0
2	C	70	0	65	1	0
All	All	6572	0	4253	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:126:SER:HA	1:A:282:TYR:HA	1.84	0.59
1:A:319:THR:O	1:A:487:GLN:NE2	2.37	0.58
1:B:369:ILE:HB	2:B:702:NAG:H81	1.89	0.53
1:A:488:LYS:NZ	1:C:508:SER:O	2.43	0.51
1:B:562:ILE:O	2:B:703:NAG:H82	2.11	0.50
1:C:382:TRP:CZ3	1:C:384:PRO:HG3	2.49	0.48
1:A:539:ARG:NH1	1:A:543:ARG:O	2.49	0.45
1:B:297:LEU:HB2	1:B:340:PHE:CD2	2.53	0.44
1:C:76:ASN:OD1	2:C:703:NAG:N2	2.54	0.40
1:C:542:MET:SD	1:C:605:TYR:CG	3.15	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	422/648 (65%)	412 (98%)	10 (2%)	0	100	100
1	B	296/648 (46%)	288 (97%)	8 (3%)	0	100	100
1	C	329/648 (51%)	321 (98%)	8 (2%)	0	100	100
All	All	1047/1944 (54%)	1021 (98%)	26 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	127/573 (22%)	127 (100%)	0	100	100
1	B	45/573 (8%)	44 (98%)	1 (2%)	45	64
1	C	98/573 (17%)	98 (100%)	0	100	100
All	All	270/1719 (16%)	269 (100%)	1 (0%)	81	83

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	563	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	487	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	NAG	A	703	1	14,14,15	1.15	1 (7%)	17,19,21	1.23	2 (11%)
2	NAG	C	705	1	14,14,15	1.14	1 (7%)	17,19,21	1.11	2 (11%)
2	NAG	A	701	1	14,14,15	1.10	1 (7%)	17,19,21	1.16	2 (11%)
2	NAG	C	701	1	14,14,15	1.09	1 (7%)	17,19,21	1.13	2 (11%)
2	NAG	A	704	1	14,14,15	1.11	1 (7%)	17,19,21	1.19	2 (11%)
2	NAG	B	701	1	14,14,15	1.12	1 (7%)	17,19,21	1.16	2 (11%)
2	NAG	B	702	1	14,14,15	1.22	1 (7%)	17,19,21	1.02	2 (11%)
2	NAG	A	702	1	14,14,15	1.14	1 (7%)	17,19,21	1.16	2 (11%)
2	NAG	C	703	1	14,14,15	1.21	1 (7%)	17,19,21	1.07	2 (11%)
2	NAG	C	704	1	14,14,15	1.11	1 (7%)	17,19,21	1.18	2 (11%)
2	NAG	A	706	1	14,14,15	1.12	1 (7%)	17,19,21	1.20	2 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\|>2$	Counts	RMSZ	$\#\|Z\|>2$
2	NAG	C	702	1	14,14,15	1.10	1 (7%)	17,19,21	1.11	2 (11%)
2	NAG	A	705	1	14,14,15	1.08	1 (7%)	17,19,21	1.15	2 (11%)
2	NAG	B	703	1	14,14,15	1.15	1 (7%)	17,19,21	1.19	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	703	1	-	0/6/23/26	0/1/1/1
2	NAG	C	705	1	-	0/6/23/26	0/1/1/1
2	NAG	A	701	1	-	0/6/23/26	0/1/1/1
2	NAG	C	701	1	-	0/6/23/26	0/1/1/1
2	NAG	A	704	1	-	0/6/23/26	0/1/1/1
2	NAG	B	701	1	-	0/6/23/26	0/1/1/1
2	NAG	B	702	1	-	2/6/23/26	0/1/1/1
2	NAG	A	702	1	-	0/6/23/26	0/1/1/1
2	NAG	C	703	1	-	0/6/23/26	0/1/1/1
2	NAG	C	704	1	-	1/6/23/26	0/1/1/1
2	NAG	A	706	1	-	3/6/23/26	0/1/1/1
2	NAG	C	702	1	-	0/6/23/26	0/1/1/1
2	NAG	A	705	1	-	2/6/23/26	0/1/1/1
2	NAG	B	703	1	-	0/6/23/26	0/1/1/1

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	703	NAG	C1-C2	3.92	1.57	1.52
2	B	702	NAG	C1-C2	3.58	1.57	1.52
2	A	703	NAG	C1-C2	3.51	1.57	1.52
2	C	705	NAG	C1-C2	3.49	1.57	1.52
2	C	704	NAG	C1-C2	3.39	1.57	1.52
2	B	703	NAG	C1-C2	3.38	1.57	1.52
2	A	706	NAG	C1-C2	3.36	1.56	1.52
2	C	702	NAG	C1-C2	3.34	1.56	1.52
2	B	701	NAG	C1-C2	3.32	1.56	1.52
2	A	702	NAG	C1-C2	3.32	1.56	1.52
2	A	701	NAG	C1-C2	3.29	1.56	1.52
2	C	701	NAG	C1-C2	3.28	1.56	1.52
2	A	704	NAG	C1-C2	3.26	1.56	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	705	NAG	C1-C2	3.21	1.56	1.52

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	703	NAG	C8-C7-N2	2.83	120.82	116.12
2	A	706	NAG	C8-C7-N2	2.75	120.68	116.12
2	B	703	NAG	C8-C7-N2	2.74	120.66	116.12
2	A	702	NAG	C8-C7-N2	2.67	120.55	116.12
2	A	704	NAG	C8-C7-N2	2.67	120.54	116.12
2	C	701	NAG	C8-C7-N2	2.65	120.51	116.12
2	A	705	NAG	C8-C7-N2	2.59	120.42	116.12
2	A	706	NAG	C2-N2-C7	-2.59	119.43	122.90
2	B	701	NAG	C8-C7-N2	2.59	120.41	116.12
2	A	701	NAG	C8-C7-N2	2.58	120.39	116.12
2	A	703	NAG	C2-N2-C7	-2.56	119.47	122.90
2	C	704	NAG	C8-C7-N2	2.56	120.36	116.12
2	C	702	NAG	C8-C7-N2	2.52	120.30	116.12
2	C	705	NAG	C8-C7-N2	2.51	120.29	116.12
2	C	703	NAG	C8-C7-N2	2.50	120.26	116.12
2	C	701	NAG	C2-N2-C7	-2.49	119.56	122.90
2	B	703	NAG	C2-N2-C7	-2.44	119.63	122.90
2	A	704	NAG	C2-N2-C7	-2.43	119.65	122.90
2	A	702	NAG	C2-N2-C7	-2.29	119.84	122.90
2	B	701	NAG	C2-N2-C7	-2.28	119.84	122.90
2	A	705	NAG	C2-N2-C7	-2.27	119.86	122.90
2	A	701	NAG	C2-N2-C7	-2.26	119.87	122.90
2	B	702	NAG	C8-C7-N2	2.22	119.80	116.12
2	C	704	NAG	C2-N2-C7	-2.21	119.93	122.90
2	C	702	NAG	C2-N2-C7	-2.14	120.03	122.90
2	B	702	NAG	C4-C3-C2	-2.13	107.90	111.02
2	C	705	NAG	C2-N2-C7	-2.09	120.10	122.90
2	C	703	NAG	C2-N2-C7	-2.02	120.19	122.90

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	705	NAG	O5-C5-C6-O6
2	A	706	NAG	C4-C5-C6-O6
2	A	706	NAG	O5-C5-C6-O6
2	C	704	NAG	O5-C5-C6-O6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	702	NAG	C1-C2-N2-C7
2	A	705	NAG	C4-C5-C6-O6
2	B	702	NAG	C3-C2-N2-C7
2	A	706	NAG	C1-C2-N2-C7

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	702	NAG	1	0
2	C	703	NAG	1	0
2	B	703	NAG	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.


6 Map visualisation

This section contains visualisations of the EMDB entry EMD-74807. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)


6.2.1 Primary map



X Index: 200




Y Index: 200




Z Index: 200


6.2.2 Raw map



X Index: 200



Y Index: 200



Z Index: 200


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)


6.3.1 Primary map



X Index: 189



Y Index: 186




Z Index: 198


6.3.2 Raw map



X Index: 189



Y Index: 186




Z Index: 198


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) (i)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views (i)


6.5.1 Primary map



X Y Z

The images above show the 3D surface view of the map at the recommended contour level 0.3. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



X Y Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate



The volume at the recommended contour level is 42 nm^3; this corresponds to an approximate mass of 38 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum (i)




*Reported resolution corresponds to spatial frequency of 0.244 Å^{-1}

8 Fourier-Shell correlation

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC



*Reported resolution corresponds to spatial frequency of 0.244 Å^{-1}

8.2 Resolution estimates (i)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.10	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	4.15	4.42	4.18

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.


9 Map-model fit

This section contains information regarding the fit between EMDB map EMD-74807 and PDB model 9ZUD. Per-residue inclusion information can be found in section 3 on page 5.


9.1 Map-model overlay



X




Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.3 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.


9.2 Q-score mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.3).

9.4 Atom inclusion (i)



At the recommended contour level, 59% of all backbone atoms, 53% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.3) and Q-score for the entire model and for each chain.



Chain	Atom inclusion	Q-score
All	0.5260	0.3090
A	0.5450	0.3180
B	0.3530	0.2540
C	0.6400	0.3410